

Author's Final Response

We thank the reviewers and the editor for helpful comments on the paper.

We have reconsidered and shortened the title as requested by reviewer 1, in line with the companion paper. Below we consider major points raised by the reviewers.

(I) Organisation of the paper (issue 1 from Reviewer 2)

We reorganized parts of the paper to give it a clearer line of thought. The distribution of SOC is based on the revised HWSD only and not on the overall 'best estimate' because the sources used to derive the best estimate could only be joined with the HWSD at global level but not at 0.5 arc minute resolution. Therefore, we first present an overview of spatial databases (maps), then present the spatial distribution based on the HWSD and derive improved global estimates from 'best estimates' of additional sources.

Within the *Introduction* we have moved up the paragraph dealing with the uncertainty of calculating C-org stocks to prepare the ground for our adjustments of bulk density. Furthermore, we moved the description of the ISRIC-WISE soil profile database from the *Results* to the *Introduction*. The *Introduction* is now followed by a section a "Spatial databases of global soil organic carbon mass". Joining the descriptions of the HWSD and other databases which were spread over the *Methods* and *Results/Discussion* sections aligns the paper with the intended review style. We removed the WRB section as no global SOC mass is derived from it. The *Results/Discussion* section is now renamed "Spatial distribution of SOC mass". We further moved several other, shorter parts of text to improve the structure or the text.

(II) Originality of the paper (Reviewer 1)

Reviewer 1 mentioned the history of global estimates of SOC. Our paper builds on this history and points at a shortcoming of a frequently used reference database, the HWSD. Our paper stands out from earlier accounts of global estimates in presenting frequency distributions of SOC that can be used by modellers for validation/calibration, focusing efforts of conserving SOC, and estimating uncertainty in studies where SOC variability was not considered.

(III) Status of the paper as a review paper

Our paper contains clearly aspects of both review and original research, but we believe the review part to dominate. It is the function of a review paper to revise assessments using existing data and to combine it with data from additional sources. Both reviewers pointed at the merit of the review, especially the unique combination of different sources. The eye-catching differences in global soil mass caused by different estimates of BD of organic soil will hopefully spur better estimates and more in-field measurements of BD.

We are looking forward to further comments by the reviewers.

| Anonymous Referee #1 | Authors' comments |
|--|---|
| <p>Soils stores a large fraction of terrestrial organic carbon. The soil organic carbon plays an important role in the biosphere. A reliable data set is the prerequisite to accurately represent the pools and fluxes of carbon. There are several estimation works based on global soil mapping (observation derived) and earth system models (simulation), and given the terrestrial organic carbon in a large range from 500 to 3000 Pg or higher. This manuscript gave us another number and geographic distribution. It is also valuable to improve our understanding on carbon cycle as a reference or benchmark data set. This manuscript gave us another estimation of soil carbon based on HWSD with adjusting the bulk density of Histosols, the definition of wetland, and incorporating more detailed estimates for permafrost from the Northern Circumpolar Soil Carbon Data Base. Though there is no anything original, it still is an important approach and valuable, and could be accepted after revision.</p> | <p>Thank you very much for taking the time to review our paper.</p> |
| <p>General comments: Soil depth matters in the SOC stock estimate, especially for deep soils. Though this paper corrected SOC of depth >1m in the peatlands, most of the true soil depth are not known in soil profile observations. Though we may lack data or method (extrapolation?) to reduce the uncertainty caused by the assumed soil depth, we need to keep this in mind. This uncertainty is not only for the <1m soil but also especially for the deeper soils. As a result, the uncertainty should be emphasized in some parts of the paper, such as page 338, line 8-17 and the conclusion section.</p> | <p>We agree that SOC stocks of deep soils are associated with great uncertainty. We emphasize in our paper the stock in the top 1 m and associated uncertainties. Stocks of deeper soils are mentioned to complement existing estimates but are not our main objective. We point out the associated uncertainty (p 338, ll 10-17, p 343, l14) and will additionally be mentioned at the end of section 3.5 in the revised text.</p> |
| <p>Consider describe the correction of frozen soils ..., and tropic peatland (Page et al.) in the method section, since the combination of the three dataset is more reliable.</p> | <p>Although the correction makes the estimate of global SOC mass more complete, it is not spatially explicit and would not allow the calculation of percentiles and masses within categories. We intended to reduce confusion by reporting in the Methods section the changes applied directly to the spatially explicit HWSD and in the Results section additional corrections considering the mass.</p> |
| <p>Consider change the order of section 3.3 and 3.4, i.e. correcting first and then overlaying with the wetland data. With the above two modifications, corresponding tables and figures need to be redraw or added.</p> | <p>Both sections can be considered separately, we do not apply any corrections within either of the sections.</p> |
| <p>Consider delete “based on the Harmonized World Soil Database” in the title.</p> | <p>We will reconsider the wording of the title.</p> |
| <p>The part 2 of the paper only uses the HWSD to calibrate a SOC model, which is not very close to the part 1. It is better to treat these two parts as independent papers.</p> | <p>We will discuss the suggestion with the editor.</p> |
| <p>Specific comments: Page 328, line 15: 1325Pg, to be consistent with the number in the conclusion.</p> | <p>This will be changed.</p> |

| | |
|---|---|
| Page 330, line 6-8: WISE(v.2) was once publicly available. But it is now replaced by the WISE (V.3.1), which is available online and includes all profiles of previous versions. | Thank you for the clarification. |
| Page 331, line17-20: The description is not precise. Shangguan et al. (2014) used three soil profile database directly, i.e. China, WISE(V.3.1) and NCSS of US, and they also used estimates (produced by others) based on local soil profiles and soil maps from ESDB (Europe), SOTWISE(various regions), GSM(US), SLC (Canada) and ASRIS (Australia). | We agree, the description will be revised. |
| Page 331, line 17, line 23: change “Shangguan et al.”to“Shangguan et al.” | The typo will be corrected. |
| Page 332, line7: It should be “equal to or smaller than”. | Correct. |
| Page 332, line9-29: Almost all the soil profiles in WISE do not have a real soil depth (or depth to the bedrock or R horizon), but have the observation depth. These soils are very likely much deeper than the recording depth in WISE. Only 189 profiles in WISE have an R horizon (some have a SOC great than 0, which seem to be errors). As a result, the overestimation for Cryosols, Podisols, and Umbrisols might not happen, especially for Cryosols and Podisols. | Good point. We will include this aspect in the revision. |
| Another point is that the soil depths in both HWSD and ISRIC-WISE are an underestimation of the true soil depth in almost all cases. It should not be named as “soil depth” in the paper. You may use the term “effective” soil depth in ISRIC-WISE (v3.0), or use the term “reference soil depth” in the HWSD. | We agree and will change the wording in the revision. |
| Page 333, line7 and et al. : kg cm^{-3} | The numbers given are correct in kg/dm^{-3} . |
| Page 333, line9: Why these regressions and the R2 are different from the authors previous report, i.e. Hiederer and Kochy (2011)? They are both based on WISE3.1. The difference of BD is 0.139 for the topsoil using the regressions when OC = 12%. | Based on your comment we reviewed our documentation. The regressions in this paper are in fact based on the SPADE/M2 database. Although the different equations produce divergent global mass results, the following correction of BD of Histosols, diminishes the difference to 1Pg. This will be mentioned in the revision. |
| Page 333, line 9: $\ln(\text{C}_{\text{org}} * 100)$ | Actually, $\text{C}_{\text{org}} \cdot 10$ was used, i.e. expressed in g/100 g soil or %. This will be indicated in the revised text. |
| Page 335, line 13-17: It lacks a soil profile database with WRB classification information to develop a WRB based soil property maps. Taxonomy reference between WRB and FAO will increase the uncertainty. | We appreciate the clarification and will add the information to the text. |
| Page 335, line 23-24: the stock was estimated based on the polygon based soil map except Australia, not after rasterization. | The text will be corrected in the revision. |
| Page 336, line 23: 13.4 Mm2 | The text refers to the soil area, so 13.1 Mm^2 is correct in this sentence. |
| Page 338, line 6-7: in the other regional stocks and the stocks of soils deeper than 1 m. | The phrasing will be corrected in the revision. |

| | |
|---|---|
| <p>Table 1: The authors used some definition which is not consistent with most literatures and may bring some confusion to the readers. I suggest using the general meanings of a terminology in the literature, instead of creating some new terms. Like the following: Content: organic carbon mass/soil dry mass; ??: organic carbon mass/soil volume (I do not see any use of this term in the paper); SOC density of a layer: organic carbon mass/soil volume\timesdepth\times(1-fractionalvolume of rocks, coarse roots, and ice); SOC density of all layers:areal density of fine soil integrated over all layers to a specified depth: SOC stock: stock integrated over a specified area.</p> | <p>We expect the paper to be of use in the soil science community but also in the carbon cycle and climate modelling community. We therefore chose to use terminology that is unambiguous and close to basic physical definitions. We understand that this decision is not optimal for both scientific communities. The definition of "content" will be deleted in the revised version as it is not used.</p> |
| <p>Table 3: Tables should be self-explainable. Please explain what is the hist/soil and it is not explained in the text.</p> | <p>Hist/soil: fraction of soil area covered by Histosols. This will be added in the revised text.</p> |
| <p>Table 3: What do you want to show with so many figures in table 3, while you only mentioned the total numbers in the text? This table needs further interpretation or you may delete it.</p> | <p>The percentiles provide information about the distribution of C in different categories of permafrost. The reason for giving the percentiles is explained in section 4.2.</p> |
| <p>Table 5: It is better to show the percentage of the overlapping. Maybe use the overlap area/(GLWD +GLCC), and 50% indicate completely identical.</p> | <p>Both approaches have their merit. We present absolute values to emphasize the significance of the overlap. In the revised text we will add the areas of each category so readers can calculate the percentage of overlap if desired.</p> |

| <i>Reviewer's comments</i> | <i>Authors' response</i> |
|--|---|
| <u>General Comments</u> | |
| <p>This manuscript provides an interesting examination of different databases that can contribute to the calculation of global soil organic carbon (SOC). The challenges of estimating the extent and characteristics of both wetlands and permafrost areas are known, but the comparison of databases that attempt to address these issues nicely illustrates the current situation. The authors give particular emphasis to the issue of bulk density (BD), which is a problem that deserves greater attention.</p> <p>Although much of the manuscript's content has merit, the effective communication is hindered by the text's organization. A major factor for obscuring the message is the appearance of five different points within the writing: 1) effect on SOC stock estimates from 'correcting' HWSO values for BD, 2) comparison of different databases' estimation of soil depths, 3) comparison of different databases' estimation of permafrost and wetland extents, 4) comparison of different databases' classification of wetland types, and 5) summing of global SOC stocks by latitude and wetland type. Clearly these points are related, but addressing them all in a coherent and focused matter will require careful crafting.</p> | Thank you very much for your careful reading of the manuscript. |
| <u>Specific Comments</u> | |
| <p>1. An apparent contradiction for the writing organization is the classification of this paper as a "review," but the text contains a methods section that does not describe the process for reviewing. Instead, this section describes a method for adjusting the BD in the HWSO. One possible solution for addressing this and my general concern about the paper's organization would be to use an outline similar to the following:</p> <p><i>I. Intro - setup of the problem, definition of key terms, and a clear statement of purpose</i></p> <p><i>II. Comparison of different databases' estimation of BD</i> <i>A. add modified HWSO as an additional item of comparison</i></p> <p><i>III. Comparison of different databases' estimation of soil depth</i></p> <p><i>IV. Comparison of different databases' estimation of permafrost and wetland extents</i> <i>A. sub-discussion on the different wetland classifications used and impact on results</i></p> <p><i>V. Summarize/compile SOC mass (summed stocks) following predefined lines of data sources by wetland category and spatially on a map. Then compare the final results of these different calculation pathways.</i></p> | We have reorganized the text by moving parts of the Methods to the Introduction and re-arranging sections within the Methods. |
| <p>2. Terms and abbreviations need to be used consistently, e.g. 0.5 arc minute v. 0.5', harmonization v. harmonisation (both acceptable spellings, choose one), SOC stocks v. organic C stocks v. organic carbon stocks.</p> | This will be addressed in the revision. |
| <p>3. P 326, L 3-6 - This needs elaborated on. Specifically, what constitutes 'relevant'?</p> | We recast the introduction so that our intention becomes clearer. |

| | |
|--|---|
| <p>4. P 332, L 8 - Is it really fair to say that the SOC stock is not underestimated with a reference soil depth of 100 cm? There are several studies showing notable amounts of SOC below 1 m (e.g. Richter and Markewitz, 1995, among others). Both in this manuscript and the published literature the qualifier of “SOC stock in the upper 1 m” is often used, which is an important distinction for what is actually being estimated. Also, later in the manuscript estimations of SOC for depths below 1 m are discussed. The subsequent breakdown of soil depths by soil type is interesting, but I suspect there is a disconnect between the definitions of sampling depth, soil depth, and the depth at which organic carbon can be found. Consideration of these issues should be part of this discussion.</p> | <p>We concur that there are considerable amounts of SOC in greater depths than 1 m. We will phrase the text more exactly in the revised text in this section.</p> |
| <p>5. P 332, L 19-22 - These sentences appear contradictory. If WISE and HWSO give the same soil depth for 80% of the area and WISE gives less soil depth for the remaining 19%, how does it work out that in total WISE gives greater depth?</p> | <p>One of the analyses was based on the WISE gridded data set which uses a maximum reference depth of 1 m. We will remove the discussion of differences in soil depth between the databases in the revised text as it goes beyond the scope of the paper.</p> |
| <p>6. P 333, L 4-6 - Provide the original HWSO 1.1 Pg C calculation as a baseline.</p> | <p>We have added a new Table 2 in the revised text.</p> |
| <p>7. P 333, L 16 - Should “mean” be inserted before “BD”?</p> | <p>It's the "best estimate" provided by Page et al., this will be clarified in the revised text.</p> |
| <p>8. P 333, L 24-27 - The difference between 2476 Pg and 1062 Pg (1414 Pg or more than 50%) does not sound “small,” but the intended comparison is probably with the 1061 Pg of the modified HWSO 1.1 calculation. Please clarify.</p> | <p>Correct. This will be clarified in the revision.</p> |
| <p>9. The comparisons of numbers are often difficult to follow. Better organization could help this, but the text at times needs to be more clear about to which number a new calculation is being compared. Tables may be helpful for this.</p> | <p>This will be addressed by the reorganization of the text.</p> |
| <p>10. P 336, L 18-20 describes the importance of the spatial mapping’s quality for frozen high-latitude soils, but only the attribute accuracy is identified as important for the global carbon mass. The area of an applied attribute is a major multiplier in any calculation of total mass. Some balance is needed to communicate that both spatial and attribute accuracy is important, but different aspects are more of a problem for the current mapping of SOC in certain land use types.</p> | <p>We agree. We will revise the text in this section and the introduction to emphasize this point.</p> |
| <p>11. P 337, L 1-2 - It appears that the CAMP map is not identifying a separate region, but a unique delineation encompassing many of the same areas as the others. If that is the case, then “a third permafrost region” should be changed to “a third permafrost extent.”</p> | <p>Correct.</p> |
| <p>12. P 339, L 19 - Is this calculation really based on an “intersection” of the two databases or the ‘union’ of the two? An intersection would be a conservative estimate, but a union seems likely to be closer to reality.</p> | <p>Here (and in line 23) it is an intersection, i.e. the area of the HWSO that is also classified as wetland.</p> |

1 Global distribution of soil organic carbon — Part 1: Masses and
2 frequency distributions of SOC stocks for the tropics,
3 permafrost regions, wetlands, and the world.

Martin Köchy 2015-1-16 10:13

Gelöscht: Soil

Martin Köchy 2015-1-16 10:13

Gelöscht: masses

Martin Köchy 2015-1-16 10:13

Gelöscht: based on the Harmonized World Soil Database

5 Martin KÖCHY^{1,a}, Roland HIEDERER², Annette FREIBAUER³

7 [1] {Thünen Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig,
8 Germany, Germany}

Martin Köchy 2015-1-16 10:13

Gelöscht:)(

Martin Köchy 2015-1-16 10:13

Gelöscht: Market Analysis)

9 [2] {Joint Research Centre, Institute for Environment and Sustainability, Via E. Fermi 2749,
10 21027 Ispra (VA), Italy}

Martin Köchy 2015-1-16 10:13

Gelöscht:)(

Martin Köchy 2015-1-16 10:13

Gelöscht:)

11 [3] {Thünen Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig,
12 Germany}

Martin Köchy 2015-1-16 10:13

Gelöscht:)(

Martin Köchy 2015-1-16 10:13

Gelöscht:)

13 [a] {now at: Thünen Institute of Market Analysis, Bundesallee 50, 38116 Braunschweig,
14 Germany}

16 Correspondence to: Martin Köchy (office@martinkoechy.de)

17

28 **Abstract**

29 The global soil organic carbon (SOC) mass is relevant for the carbon cycle budget and thus
30 atmospheric carbon concentrations. We review current estimates of SOC stocks and mass
31 (stock × area) in wetlands, permafrost and tropical regions and the world in the upper 1 m of
32 soil. The Harmonized World Soil Database (HWSD) v.1.2 provides one of the most recent
33 and coherent global data sets of SOC, giving a total mass of 2476 Pg when using the original
34 values for bulk density. Adjusting the HWSD’s bulk density of soil high in organic carbon
35 results in a mass of 1230 Pg, and additionally setting the BD of Histosols to 0.1 g cm⁻³
36 (typical of peat soils) results in a mass of 1062 Pg. The uncertainty of bulk density of
37 Histosols alone introduces a range of –56 to +180 Pg C for the estimate of global SOC mass
38 in the top 1 meter, larger than estimates of global soil respiration. We report the spatial
39 distribution of SOC stocks per 0.5 arc minutes, the areal masses of SOC and the quantiles of
40 SOC stocks by continents, wetland types, and permafrost types. Depending on the definition
41 of ‘wetland’, wetland soils contain between 82 and 158 Pg SOC. Incorporating more detailed
42 estimates for permafrost from the Northern Circumpolar Soil Carbon Data Base (496 Pg
43 SOC) and tropical peatland carbon, global soils contain 1325 Pg SOC in the upper 1 m
44 including 421 Pg in tropical soils, whereof 40 Pg occur in tropical wetlands. Global SOC
45 amounts to just under 3000 Pg when estimates for deeper soil layers are included. Variability
46 in estimates is due to variation in definitions of soil units, differences in soil property
47 databases, scarcity of information about soil carbon at depths >1 m in peatlands, and variation
48 in definitions of ‘peatland’.

Martin Köchy 2015-1-16 10:13

Gelösch:

Martin Köchy 2015-1-16 10:13

Gelösch: soil organic carbon

Martin Köchy 2015-1-16 10:13

Gelösch: (mass/area)

Martin Köchy 2015-1-16 10:13

Gelösch: . Correcting

Martin Köchy 2015-1-16 10:13

Gelösch: soils, especially

Martin Köchy 2015-1-16 10:13

Gelösch: .

Martin Köchy 2015-1-16 10:13

Gelösch: 1324

56 **1. Introduction**

57 The global mass of soil organic carbon (SOC) is greater than the combined mass of carbon
58 (C) contained in the atmosphere and in the living biomass. Therefore, small changes in the
59 mass of SOC can have profound effects on the concentration of atmospheric CO₂ and hence
60 climate change. Despite its importance, the global mass of SOC and its distribution in space
61 and among land-use/land-cover classes is not well known.

62 On the short to middle term (decades), variation in SOC mass is strongly related to the
63 balance of input from net primary productivity and microbial decomposition. On longer time-
64 scales, however, changes in the decomposable mass of SOC affect this balance. Globally, the
65 largest SOC stocks are located in wetlands and peatlands, most of which occur in regions of
66 permafrost and in the tropics. Decomposition rates in wetlands and permafrost are low due to
67 low availability of oxygen and low temperatures, respectively. This SOC is vulnerable to
68 changes in the hydrological cycle as well as to changes in permafrost dynamics.

69 A good knowledge of the global SOC mass and its spatial distribution is necessary for
70 assessing, in an international context, where soils are most vulnerable to C losses or which
71 land use/land cover types might provide the best opportunity for C sequestration to mitigate
72 increases in greenhouse gas concentrations. Since SOC mass is a product of several factors,
73 uncertainty (or errors in measurement) in one of the factors affects all others. Consequently,
74 the measures to reduce the uncertainty of global SOC mass should be directed to those soils
75 that are associated with a large extent (area), high levels of C_{org}, low bulk density (BD) or
76 great depth. Variations at the lower end of BD are more consequential than at the high end of
77 BD because low BD is associated with organic soils (high C_{org}) and a change from, say, 0.1 to
78 0.2 leads to a doubling of SOC stock and mass. Variation within the range of BD typical of
79 mineral soils, 1.2 – 1.8 g cm⁻³ is less consequential.

Martin Köchy 2015-1-16 10:13

Gelösch: The large terrestrial organic carbon stocks of soil and biomass interact closely.

Martin Köchy 2015-1-16 10:13

Gelösch: more

Martin Köchy 2015-1-16 10:13

Gelösch: , but on

Martin Köchy 2015-1-16 10:13

Gelösch: become more relevant

86 Traditionally, maps of the spatial distribution of SOC stocks are derived from maps where
87 areas with similar soil characteristics are aggregated to form soil mapping units, and the SOC
88 mass of the area of the soil mapping unit is calculated by multiplication of the area of the soil
89 mapping unit with its unit-area SOC stock (Amundson, 2001). Historically, soil maps have
90 been compiled largely based on the experience of soil surveyors, taking into account
91 topography, climate, land use history, land management, vegetation, parent material, and soil
92 typical characteristics (McBratney et al., 2003b). The spatial soil mapping units are linked to
93 their defining properties, which are based on measurements of soil profiles or an evaluation
94 by experts. Typically, measurements from several profiles within the same soil unit have been
95 statistically aggregated (e.g., averaged). Missing profile data may be estimated by
96 pedotransfer functions (PTF) from other measured soil characteristics.

97 The SOC stock, m_C , of a soil column is calculated by integrating the areal density of SOC
98 over all vertical depth layers (or within a specified depth). The areal density of SOC of a soil
99 layer is determined by measuring the organic carbon concentration (C_{org}) and the bulk density
100 (BD) of undisturbed soil samples in homogenous layers of thickness d (Table 1). The areal
101 density, $C_{org} \times BD \times d$, is reduced by the fractional volume f_G occupied by gravel, rocks, roots,
102 and ice in the soil layer, or $m_C = C_{org} \times BD \times (1 - f_G) \times d$. The SOC mass of the area (A) is the
103 product of the soil unit's area and its SOC stock ($m_C \times A$). Lateral variability, temporal
104 variability, and methodological differences in measuring any of the necessary soil
105 characteristics (BD, C_{org} , volume of gravel and roots, forms of C, depth) contribute to the
106 variability of SOC stock and mass estimates (Ellert et al., 2001).

107 The accuracy of spatially interpolated maps of SOC stocks depends on how well the soil
108 mapping units are represented by soil profiles with complete characteristics. The latest ISRIC-
109 WISE database (v.3.1) contains harmonized data of more than 10250 soil profiles (Batjes,
110 2009). The profiles, however, do not yet represent the terrestrial surface equally. Gaps include

Martin Köchy 2015-1-16 10:13

Gelösch: stock per area (henceforth SOC)

Martin Köchy 2015-1-16 10:13

Gelösch:)

Martin Köchy 2015-1-16 10:13

Gelösch: summation over

Martin Köchy 2015-1-16 10:13

Gelösch: underlying

Martin Köchy 2015-1-16 10:13

Gelösch: that have been classified as the same soil unit.

Martin Köchy 2015-1-16 10:13

Gelösch: on

Martin Köchy 2015-1-16 10:13

[1] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13

Gelösch: A good knowledge of the global SOC mass and its spatial distribution is necessary for assessing in an international context which soils are most vulnerable or might provide the best opportunity for C sequestration in mitigation of rising greenhouse gas concentrations. At the global scale, in-situ measurements must be complemented by modelling activities, which are greatly improved if variation in key factors like soil organic carbon can be accounted for. In this paper we review existing spatial estimates of SOC stocks and masses, including their uncertainties and underlying methods for estimating the stocks. Our paper reports for the first time area-weighted frequency distributions of carbon stocks within land-use and land-cover classes, using best estimates from several sources. We focus on the large SOC stocks in wetlands, tropical soil, and permafrost at high latitudes and present frequency distributions of SOC stocks within classes of land use, land cover, and geographic region. Furthermore, we provide recommendations for improving global soil mapping. .

... [1]

145 non-agricultural areas of North America, the Nordic countries, most parts of Asia (notably
146 Iran, Kazakhstan, and Russia), Northern Africa, and Australia. To calculate SOC stocks one
147 needs C_{org} , BD, soil depth, and volumetric gravel fraction. These are provided individually by
148 87%, 32%, 100%, and 22%, respectively, of the profiles (Batjes, 2009). BD and gravel
149 fraction have low representation because they are seldom recorded during routine soil
150 surveys. In numbers, 9970 profile descriptions include C_{org} in at least one layer, but of these
151 only 3655 also include BD. Gravel fraction is explicitly indicated for 1100 of the 3655
152 profiles but earlier versions of the database could not distinguish between zero and absence of
153 value. BD is included for 806 profiles where $C_{org} > 3\%$ and for 74 profiles where $C_{org} > 20\%$.
154 The temporal origin of profile descriptions ranges from 1925 to 2005. The early data may no
155 longer reflect current conditions where C input and decomposition rates may have changed.
156 Efforts to expand the database of data-rich soil profiles and to use pedotransfer instead of
157 taxotransfer functions has been going on since 1986 through the SOTER program
158 (<http://www.isric.org/projects/soil-and-terrain-database-soter-programme>, accessed 2014-07-
159 07, Nachtergaele, 1999).

160 In this paper we compare previous estimates of the global SOC mass in the top 1 m of soil
161 derived from spatial databases (maps) to the mass derived from the Harmonized World Soil
162 Database (HWSD, FAO et al., 2012), which was the latest and most detailed inventory at the
163 global scale when this study was begun and is still widely used as an international reference
164 (e.g., Wieder et al., 2014, Yan et al., 2014) but requires adjustment of bulk densities of
165 organic soils (Hiederer & Köchy 2011). Based on the adjusted HWSD, our paper reports for
166 the first time area-weighted frequency distributions of carbon stocks in the top 1m of soil, in
167 particular for the large SOC stocks in wetlands, the tropics, and permafrost at high latitudes.
168 Frequency distributions can be used to improve the assessment of accuracy in studies where
169 SOC is an independent variable. We update the HWSD-derived global SOC mass using best

Martin Köchy 2015-1-16 10:13
[2] verschoben (Einfügung)

170 [estimates for the permafrost region, the tropics, and soil below 1 m depth from several](#)
171 [additional sources. Our conclusions provide recommendations for improving global soil](#)
172 [mapping.](#)

173 **2. Spatial databases of global soil organic carbon mass**

174 Historic estimates of global SOC mass compared among 27 studies range between 504 and
175 3000 Pg with a median of 1461 Pg (Scharlemann et al., 2014). Here we concentrate on
176 comparisons with the most recent ones.

177 The Harmonized World Soil Database (HWSD vers. 1.2, FAO et al., 2012) with a raster of
178 0.5 arc minutes (0.5') is one of the most recent and most detailed databases at the global scale
179 and widely used as reference. The HWSD contains for the topsoil (0-30 cm) and the subsoil
180 (30-100 cm) values for C_{org}, BD and gravel content for dominant and secondary soil types on
181 a 0.5' grid. Data sources for HWSD are earlier global soil maps that were published by or in
182 cooperation with FAO, the European Soil Database, the Soil Map of China, SOTER regional
183 studies, WISE profile data, and WISE pedotransfer and taxotransfer functions. The HWSD
184 does not yet include the extensive national databases of USA, Canada, and Australia. The
185 HWSD is the result of associating existing maps of soil types (if necessary reclassified to
186 FAO standards) with soil characteristics derived from the WISE (v.2) database containing
187 about 9600 soil profiles, which is the largest number used for a global soil map until 2013.

188 The HWSD does not quantify variability or ranges of any soil properties within a soil unit. Its
189 description qualifies that “Reliability of the information contained in the database is variable:
190 the parts of the database that still make use of the Soil Map of the World such as North
191 America, Australia, West Africa and South Asia are considered less reliable, while most of
192 the areas covered by SOTER databases are considered to have the highest reliability (Central
193 and Southern Africa, Latin America and the Caribbean, Central and Eastern Europe).”

194 A default reference soil depth of 100 cm is stipulated in the HWSD for each mapping unit as a
195 concession to harmonization of different soil databases. Only Rendzinas, Rankers, Leptosols,
196 and Lithosols are attributed reference soil depths of 30 cm or 10 cm. For most of the

Martin Köchy 2015-1-16 10:13
[3] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13

Gelöscht: 1

Martin Köchy 2015-1-16 10:13

Gelöscht: 2009

Martin Köchy 2015-1-16 10:13

Gelöscht: because it was

Martin Köchy 2015-1-16 10:13

Gelöscht: latest

Martin Köchy 2015-1-16 10:13

Gelöscht: inventory

Martin Köchy 2015-1-16 10:13

Gelöscht: when this study was begun. The database was updated to version 1.2.1 in March 2012 with minor effects on the results presented here (details follow below).

Martin Köchy 2015-1-16 10:13

Gelöscht: 5 arc minute grid. Data derived from HWSD and co-published with HWSD (Fischer et al., 2008) include O₂ constraint and presence of permafrost at 5 arc minute resolution.

Martin Köchy 2015-1-16 10:13

Gelöscht: (v.1.2)

Martin Köchy 2015-1-16 10:13

Gelöscht: but is not available publicly. A short account of earlier published maps of SOC is presented in the Results.

215 remaining soil units the 25-percentile of lowest recorded depth of profiles in the WISE 3.1
216 database is equal to or greater than the reference depth, i.e., SOC stock within the top 1 m is
217 not underestimated by using the reference depth. The 25-percentiles of recorded depths of
218 Calcisols (95 cm, n=218), Cambisols (90 cm, n=1164), Cryosols (80 cm, n=6), Durisols (45
219 cm, n=1), Podisols (80 cm, n=222), Solonchaks (90 cm, n=165), and Umbrisols (49 cm,
220 n=173) are smaller than the reference depths so that C stocks may be overestimated. The
221 overestimate could be substantial for Cryosols, Podisols, and Umbrisols, which have high C_{org}
222 (median >10%). At the same time, the true soil depth of Cryosols and Podisols can be
223 expected to be deeper than the recorded depth in the databases, which, however, would be of
224 no consequence for the estimated SOC mass of the top 1 m.

225 The global SOC mass calculated directly from the original HWSD (v.1.1) for the upper 1 m
226 of soil is 2476 Pg, Henry *et al.* (2009), using an unspecified earlier version of HWSD,
227 reported a mass of 1850 Pg for the first meter. These high values are, however, due to
228 inconsistencies, gaps, and inaccuracies in the database (Hiederer & Köchy, 2011). The most
229 consequential of the inaccuracies concerns the BD for soils high in C_{org}. In addressing these
230 issues (see Methods), we calculated a global mass of SOC in the top 1 m of soil of 1230 Pg in
231 a first step and 1062 Pg in a second step.

232 Before the publication of the HWSD, many global estimates were based on the Digital Soil
233 Map of the World (DSMW) (FAO, 2007) or its precursor, the Soil Map of the World (FAO,
234 1997). Batjes (1996), using information from 4353 WISE profiles, reported a range of 1462–
235 1548 Pg for 0–1 m depth and 2376–2456 Pg for 0–2 m depth, Henry *et al.* (2009) report a
236 global SOC mass of 1589 Pg for the top 1 m and 2521 Pg for the top 2 m (using an
237 unspecified WISE version). Hiederer *et al.* (2010) report a slightly lower mass of 1455 Pg for
238 DSMW for the top 1 m.

Martin Köchy 2015-1-16 10:13
[4] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13
[5] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13
[6] verschoben (Einfügung)

239 The International Geosphere Biosphere Program (IGBP) (Global Soil Data Task Group, 2000)
240 produced a map of SOC stock on a 5' by 5' grid derived from the DSMW in conjunction with
241 WISE data (v.1, 1125 profiles). SOC mass (0–1 m) based on the IGBP map is 1550 Pg
242 (calculated as SOC stock × grid cell area).

Martin Köchy 2015-1-16 10:13
[7] verschoben (Einfügung)

243 The US Natural Resources Conservation Services reclassified the SMW at 2' and combined it
244 with a soil climate map (Reich, 2000, data — on a 3' grid—, downloaded from [http://spatial-](http://spatial-analyst.net/worldmaps/SOC.zip)
245 analyst.net/worldmaps/SOC.zip). This map shows the distribution of nine classes of SOC
246 stocks that result in a global SOC mass (0–1 m) of 1463 Pg. Analyzing 2721 soil profiles
247 grouped per biome, Jobbágy & Jackson (2000) estimated that the top 1 m contains 1502 Pg
248 SOC, with 491 Pg in 1-2 m and 351 Pg in 2-3 m depth.

Martin Köchy 2015-1-16 10:13
[8] verschoben (Einfügung)

249 The recently published Global Soil Dataset for Earth System Models (Shangguan et al., 2014)
250 with a resolution of 0.5', combined the DSMW with regional soil maps and global and
251 regional profile databases from several sources beyond those used in the HWSD, including
252 the national databases of the USA, Canada, and Australia. Soil profile data and mapping units
253 were matched in several steps intended to result in the most reliable information. Several
254 harmonization steps are applied to the data to derive amongst others soil carbon
255 concentration, bulk soil density, and gravel content and depth for each soil mapping unit. The
256 global SOC stocks are reported as 1923, 1455 and 720 Pg for the upper 2.3, 1.0, and 0.3 m,
257 respectively.

Martin Köchy 2015-1-16 10:13
[9] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13
Gelöscht: Recently, Shangguan et al.

Martin Köchy 2015-1-16 10:13
Gelöscht: provided a new interpretation of the Digital Soil Map of the World (DSMW), FAO, 2007), for use in earth system modelling. They included the soil profile data from the USA, Canada, and Australia, which required additional routines of harmonization. Here, we base our analysis on the HWSD because it is still widely used as an international reference (e.g., Wieder et al., 2014, Yan et al., 2014). We also present a correction of overestimated BD values for Histosols contained in the HWSD that was not specifically addressed by Shangguan et al. (2014, further details below), Hiederer & Köchy (2011), or Scharlemann et al. (Scharlemann et al., 2014).

275 **3. Processing of HWSD data for spatial analyses**
276 We calculated the SOC stocks for each soil type (*s*) within a grid cell as the areal density over
277 the thickness of the top and sub soil layer, accounting for the volume occupied by gravel, and
278 weighted it with the soil type's areal fraction in each cell or $m_{C_s} \times A_s / \sum A$. Consequently, SOC
279 mass of each cell is the sum over all soil types of the product of SOC stock of each soil type
280 and the fraction of cell area covered by each soil type or $\sum(m_{C_s} \times A_s / A)$.

281 Our analysis of SOC stocks and masses is based on (HWSD vers. 1.1, FAO et al., 2009)
282 because it was the latest version when this study was begun. Version 1.2 of the HWSD adds
283 two new fields for BD (one for topsoil and one for subsoil) based on the SOTWIS database
284 and addresses minor issues that are listed in detail on the HWSD's web site. Since the
285 resulting differences in global mass between HWSD versions were <0.3%, we did not
286 recalculate the other values so that all values reported below are calculated based on version
287 1.1 of the HWSD and a global mass of 1061 Pg unless explicitly mentioned otherwise.

288 Despite the harmonization of the spatial and attribute data the HWSD suffers from some
289 residual inconsistencies in the data reported, gaps in some areas covered and errors in the
290 values reported (Hiederer and Köchy, 2011) that required pre-processing of the data
291 (Supplement). Here we present a correction of overestimated BD values for Histosols
292 contained in the HWSD that was not specifically addressed by Shangquan et al. (2014),
293 Hiederer & Köchy (2011), or Scharlemann et al. (Scharlemann et al., 2014). For each
294 processing step the resulting global SOC mass is used as an indication of the magnitude of the
295 data manipulation (Table 2).

Martin Köchy 2015-1-16 10:13

Gelösch: excluding

Martin Köchy 2015-1-16 10:13

Gelösch: according to

Martin Köchy 2015-1-16 10:13

Gelösch: A uniform reference soil depth of 100 cm is stipulated in the HWSD for each mapping unit as a concession to harmonization of different soil databases. Only Rendzinas, Rankers, Leptosols, and Lithosols are attributed soil depths of 30 cm or 10 cm. For most of the remaining soil units the reference depth is equal to or greater than the 25-percentile of profiles in the WISE 3.1 database, i.e., SOC stock is not underestimated by using the reference depth. The 25-percentiles of Calcisols (95 cm, n=218), Cambisols (90 cm, n=1164), Cryosols (80 cm, n=6), Durisols (45 cm, n=1), Podzols (80 cm, n=222), Solonchaks (90 cm, n=165), and Umbrisols (49 cm, n=173) are smaller than the reference depths so that C stocks may be overestimated. The overestimate might be great for Cryosols, Podzols, and Umbrisols, which have high C_{org} (median >10%). For our calculations of SOC mass reported in this paper we did not, however, correct SOC stocks for depth because it would have required a profile-by-profile check whether the recorded maximum depth may have been the end of the solum or the end of the soil sample, which was beyond the scope of this analysis. A spatial, equal-area comparison (regression) of soil depth between HWSD and the ISRIC-WISE (v3.0) 0.5 degree grid resulted in a slope coefficient of 0.82. For over 80% of the surface, WISE and HWSD give the same soil depth. For 19% of the global land surface WISE gives less soil depth than HWSD. The differences between HWSD and WISE soil depth are unevenly distributed. Globally the WISE database gives greater soil depth. Higher, but locally restricted differences are found in southern Argentina (WISE soil depth: 10cm, HWSD: 86cm). Smaller divergence, but with greater spatial cover are found mainly in China and eastern Siberia (WISE: 69cm, HWSD: 100cm). The differences in soil depth between the databases may be attributed to the different source data. With the HWSD using regional data sources one may argue that these data should better represent the regional variations. However, it was outside the scope of the study to evaluate the accuracy of the data.

Martin Köchy 2015-1-16 10:13

[10] verschoben (Einfügung)

Martin Köchy 2015-1-16 10:13

Gelösch: Appendix A).

Martin Köchy 2015-1-16 10:13

Gelösch: we report in the remainder of this paragraph

Martin Köchy 2015-1-16 10:13

Gelösch: .

353 (Step 1) We filled missing data for C_{org} in top (4 cases) and subsoil layers (127 cases) with
 354 data from cells characterized as the same soil unit and being closest in distance or most
 355 similar in topsoil C_{org} . (2) In a similar way we filled missing values of BD for mineral soils in
 356 27 cases. (3) In HWSD v.1.1 high C_{org} values (>20%) are associated with a BD of 1.1 to 1.4
 357 kg/dm^3 although values of 0.05 to 0.3 kg/dm^3 would be typical of organic soils (Boelter, 1968,
 358 Page et al., 2011). To address this issue, we set the topsoil BD to $-0.31 \ln(C_{org} [\%]) + 1.38$
 359 ($R^2=0.69$) and subsoil to $-0.32 \ln(C_{org} [\%]) + 1.38$ ($R^2=0.90$) for $C_{org} > 3\%$ based on an analysis
 360 of the SPADE/M2 soil profile database (Hiederer, 2010). This results in a global mass of
 361 1230 Pg C for a soil depth of up to 1 m. (4) The maximum C_{org} of Histosols in the HWSD is
 362 47%, resulting in a BD of 0.19 kg/dm^3 for topsoil and 0.15 kg/dm^3 for subsoil using the
 363 mentioned equations. In contrast, the best estimate for the BD for tropical peatlands is 0.09
 364 kg/dm^3 (Page et al., 2011), for boreal and subarctic peatland the average BD is 0.112 kg/dm^3
 365 (Gorham, 1991), and for Finnish agricultural peat soil the average value is 0.091 kg/dm^3
 366 (Mäkkilä, 1994 in Turunen, 2008). Therefore, we set BD to 0.1 kg/dm^3 for all Histosols in
 367 HWSD. With the SPADE/M2-based corrections for BD and the modification for Histosols,
 368 the global mass of SOC in the upper 1 m of soil is 1061 Pg. Hiederer & Köchy (2011) used
 369 WISE-based corrections for BD ($BD_{top} = -0.285 \ln(C_{org} [\%]) + 1.457$ and $BD_{sub} = -0.291$
 370 $\ln(C_{org} [\%]) + 1.389$ for $C_{org} > 12\%$) that result in a higher a global C mass of 1376 Pg in step 3
 371 but a very similar mass (1062 Pg) after the BD correction for histosols in step 4.
 372 For comparison, total SOC mass derived from the unaltered HWSD v1.2 database and using
 373 the SOTWIS BD (when available for a soil mapping unit) is 2476 Pg and 1062 Pg after
 374 applying the BD correction as described in the previous paragraph.
 375 The HWSD database was pre-processed and analyzed with R (R Development Core Team,
 376 2011). Details of the calculations are presented in the Supplement. We summarized adjusted
 377 SOC stocks from HWSD globally and by geographic regions, land cover types, and areas with

Martin Köchy 2015-1-16 10:13

Gelösch:)+

Martin Köchy 2015-1-16 10:13

Gelösch:)+

Martin Köchy 2015-1-16 10:13

Gelösch: ISRIC-WISE v3.1

Martin Köchy 2015-1-16 10:13

Gelösch: .

Martin Köchy 2015-1-16 10:13

Gelösch: Different gap-filling and modification to the BD of all organic soils result in a global C mass ranging between 1208 Pg and 1338 Pg (Carré et al., 2010, Hiederer and Köchy, 2011, Wieder et al., 2014).

Martin Köchy 2015-1-16 10:13

Gelösch:

Martin Köchy 2015-1-16 10:13

Gelösch: Based on these assumptions the global mass of organic C in the upper 1 m of soil is 1061 Pg.

Martin Köchy 2015-1-16 10:13

Gelösch: Version 1.2 of the HWSD was published after finishing the research for this paper. The new version adds two BD columns (one for top and one for subsoil) based on the SOTWIS database and addresses minor issues that are listed in detail on the HWSD's web site. Total

Martin Köchy 2015-1-16 10:13

Gelösch: using

Martin Köchy 2015-1-16 10:13

Gelösch: .

Martin Köchy 2015-1-16 10:13

Gelösch: and using the SOTWIS BD (when available for a soil mapping unit), it is 1062 Pg. Since these changes were small, we did not recalculate the other values so that all values reported below are calculated based on version 1.1 of the HWSD unless explicitly mentioned otherwise.

Martin Köchy 2015-1-16 10:13

Gelösch: analysed

Martin Köchy 2015-1-16 10:13

Gelösch: Appendix A.

410 specific soil characteristics (wetlands, peatlands, permafrost soils). To achieve this we
411 intersected raster maps of SOC with thematic maps in a GIS (GRASS 6.4.2, GRASS
412 Development Team, 2011) and calculated SOC mass summed over areas and determined the
413 5th, 25th, 50th, 75th, and 95th percentiles of SOC stocks within these areas.

414 |

Martin Köchy 2015-1-16 10:13
Gelöscht: <#>Results and Discussio ... [2]

417 **4. Spatial distribution of SOC mass**

418 **4.1. Continental distribution of SOC mass**

419 The distribution of SOC mass by continents (Table 3) follows the pattern of terrestrial

420 ecological zones. A large areal fraction of deserts obviously reduces the continental mean

421 SOC stock, whereas a large fraction of frozen organic soil increases the continental mean

422 SOC stock (Fig. 1).

423 **4.2. Carbon in frozen high-latitude soils**

424 Large SOC deposits exist in the frozen soils of the permafrost region and are vulnerable to the

425 effects of global warming. The mass of these deposits, however, is not well known because

426 area and the stocks of the permafrost region are uncertain. The uncertainty of the area is

427 characterized by the variation in the delineation and thus extent of the permafrost region

428 among different maps and databases, which is due also to different definitions of "permafrost"

429 and associated concepts. The HWSD lists for each soil unit the presence of permafrost within

430 the top 200 cm (a so-called 'gelic phase'). SOC mass in the top 1 m of soils with a gelic phase

431 is 164 Pg for a 13.1 Mm² soil area (Table 4). Supplementary data to the HWSD (Fischer et al.,

432 2008) indicate on a 5' grid the presence of continuous or discontinuous (i.e., excluding

433 sporadic and isolated) permafrost that is based on the analysis of the snow-adjusted air frost

434 number (Harrij van Velthuisen, IIASA, pers. Comm. 2011) as used for the Global Agro-

435 ecological Zones Assessment v3.0 (Fischer et al., 2008). This extent (19.5 Mm² cell area, Fig.

436 2) encompasses the area of soils with a gelic phase and contains 185 Pg SOC on 16.7 Mm²

437 soil area according to the HWSD. A third permafrost extent (24.9 Mm² cell area) is described

438 by the Circum-Arctic Map of Permafrost and Ground Ice Conditions (CAMP, Heginbottom et

439 al., 1993), which comprises 12 categories of permafrost and ground ice prevalence without a

- Martin Köchy 2015-1-16 10:13
[10] nach oben verschoben: (Scharlemann et al., 2014). ... [3]
- Martin Köchy 2015-1-16 10:13
Gelösch: The global SOC mass calculated directly from the HWSD (v.1.2.1) for the upper 1 m of soil is 2476 Pg.
- Martin Köchy 2015-1-16 10:13
[4] nach oben verschoben: Henry et al. (2009), using an unspecified earlier version of HWSD, reported a mass of 1850 Pg for the first meter. These high values are, however, due to inconsistencies, gaps, and inaccuracies in the database (
- Martin Köchy 2015-1-16 10:13
Gelösch: see Methods
- Martin Köchy 2015-1-16 10:13
[5] nach oben verschoben:). The most consequential of the inaccuracies concerns the BD for soils high in C_{org}.
- Martin Köchy 2015-1-16 10:13
Gelösch: After addressing these issues, we calculated a global mass of SOC in the top 1 m of soil to 1062 Pg. The distribution of SOC mass by continents (Table 2) follows the pattern of land area.
- Martin Köchy 2015-1-16 10:13
[6] nach oben verschoben: Henry et al. (2009) report a global SOC mass of 1589 Pg for the top 1 m and 2521 Pg for the top 2 m (using an unspecified WISE version). Hiederer et al. (2010) report a slightly lower mass of 1455 Pg for DSMW
- Martin Köchy 2015-1-16 10:13
Gelösch: and the top 1 m.
- Martin Köchy 2015-1-16 10:13
[7] nach oben verschoben: The International Geosphere Biosphere Program (IGBP) (Global Soil Data Task Group, 2000) produced a map of SOC stock on a 5' by 5' grid derived from the DSMW in conjunction with WISE data (v.1, 1125 profiles). SC... [5]
- Martin Köchy 2015-1-16 10:13
[1] nach oben verschoben: stocks depends on how well the soil mapping units are represented by soil profiles with complete characteristics. The latest ISRIC-WISE ... [7]
- Martin Köchy 2015-1-16 10:13
[2] nach oben verschoben: of the profiles (Batjes, 2009). BD and gravel f... [8]
- Martin Köchy 2015-1-16 10:13
Gelösch: Before the publication of the HWSD, many global estimates were ba... [4]
- Martin Köchy 2015-1-16 10:13
Gelösch: - ... [6]
- Martin Köchy 2015-1-16 10:13
Gelösch: These are provided individually by 87%, 32%, 100%, and 22%
- Martin Köchy 2015-1-16 10:13
Gelösch: organic C...OC deposits e... [9]

647 defined depth limit for the occurrence of permafrost. The CAMP permafrost extent (including
648 permafrost in the Alps and Central Asian ranges) represents 21.7 Mm² soil area of the HWSD
649 with 249 Pg SOC in the top 1 m. Tarnocai *et al.* (2009) used the CAMP's permafrost
650 classification (excluding the Alps and Central Asian ranges, 20.5 Mm² grid cell area) together
651 with SOC and soil information from the Northern Circumpolar Soil Carbon Data Base
652 (NCSCDB, <http://wms1.agr.gc.ca/NortherCircumpolar/northercircumpolar.zip>) to estimate
653 SOC mass in the permafrost region. The NCSCDB includes soil profile data not incorporated
654 into the HWSD. Data for calculating SOC stocks (C concentration, BD, depth, coarse
655 fragments) in the upper 3 m were derived from 1038 pedons from northern Canada, 131
656 pedons from Alaska, 253 pedons from Russia, 90 peat cores from western Siberia, 266
657 mineral and organic soils from the Usa Basin database, and an unspecified number of profiles
658 from the WISE database (v.1.1) for Eurasian soils. Extrapolations were used to estimate SOC
659 mass in mineral soils and Eurasian peat soils >1 m depth. The spatial extent of soil classes
660 was obtained from existing digital and paper maps. Tarnocai *et al.*'s (2009) estimate of 496
661 Pg for the 0–1 m depth is much higher than that of HWSD's mass in the permafrost region
662 (185 Pg). The difference is partly due the limit of 2 m that HWSD uses for distinguishing the
663 'gelic phase', whereas the Circum-arctic Map of Permafrost does not refer to a limit
664 (Heginbottom *et al.*, 1993). The more important contribution to the difference in mass than
665 arising from definitions and extent is the greater SOC stock calculated from the NCSCDB
666 (Table 4). In NCSCDB the mean SOC areal density of soil in all permafrost classes is >20
667 kg/m², whereas the mean SOC areal density is 11.4 kg/m² in the HWSD across all classes.
668 The difference suggests that the BD of frozen organic soil is higher than assumed by us. In
669 addition to the SOC mass in the top 1 m, Tarnocai *et al.* (2009) estimated that the permafrost
670 region contains 528 Pg in 1 m to 3 m depth, and 648 Pg in depths greater than 3 m.

Martin Köchy 2015-1-16 10:13

Gelösch: area comprises 21.7 Mm² soil area of the HWSD

Martin Köchy 2015-1-16 10:13

Gelösch: cause of

Martin Köchy 2015-1-16 10:13

Gelösch: stock

Martin Köchy 2015-1-16 10:13

Gelösch: stock

676 Inaccuracies associated with the mass estimates arise from incomplete knowledge of the
677 spatial distribution of soil classes, soil depths, sparse distribution of soil profile data and lack
678 of soil profiles with a full complement of measured data. In terms of IPCC A4 categories of
679 confidence, Tarnocai *et al.* have medium to high confidence (>66%) in the values for the
680 North-American stocks of the top 1 m, medium confidence (33–66%) in the values for the
681 Eurasian stocks of the top 1 m, and very low to low confidence (<33%) in the values for the
682 other regional stocks and stocks of layers deeper than 1 m. Tarnocai *et al.* discuss extensively
683 the uncertainty of their estimates. Here we note only that major uncertainty is linked to the
684 area covered by high latitude peatlands (published estimates vary between 1.2 and 2.7 Mm²)
685 which alone results in a range of 94–215 Pg SOC. The C mass contained in >3 m depth of
686 river deltas is potentially great (241 Pg, Tarnocai *et al.*, 2009), but is based solely on
687 extrapolation on the SOC stock and area of the Mackenzie River delta. Yedoma (Pleistocene
688 loess deposits with high C_{org}) SOC mass (407 Pg, >3 m depth) is also associated with great
689 uncertainty. The estimate (adopted from Zimov *et al.*, 2006) is based on a sketched area of 1
690 Mm² in Siberia (thus excluding smaller Yedoma deposits in North America) and mean
691 literature values for depth (25 m) whose ranges extend >±50% of the mean.

692 4.3. Carbon in global wetlands

693 SOC stocks in wetlands are considerable because water reduces the availability of oxygen and
694 thus greatly reduces decomposition rates (Freeman *et al.*, 2001). Draining of wetlands often
695 greatly increases the decomposition of dead plant material, which results in the release of
696 carbon dioxide into the atmosphere. This process can significantly affect the global C budget
697 when it happens on a large scale. There is, however, no consensus of what constitutes a
698 wetland at the global scale (Mitra *et al.*, 2005). Therefore the volume of wetland soil and its C
699 mass are also uncertain (Joosten, 2010).

Martin Köchy 2015-1-16 10:13

Gelösch: mass

Martin Köchy 2015-1-16 10:13

Gelösch: mass

Martin Köchy 2015-1-16 10:13

Gelösch: masses

703 The most detailed and recent maps of global scope with detailed wetland classification
704 (Köchy and Freibauer, 2009) are the Global Land Cover Characteristics database, v 2.0
705 (GLCC, Loveland et al., 2000) that comprises up to 6 wetland types ('Wooded Wet Swamp',
706 'Rice Paddy and Field', 'Inland Water', 'Mangrove', 'Mire, Bog, Fen', 'Marsh Wetland') and
707 the Global Lakes and Wetland Database (GLWD, Lehner and Döll, 2004) that comprises 12
708 wetland categories. Both maps have a resolution of 0.5'. The GLCC originates from analysis
709 of remote sensing data in the International Geosphere Biosphere program. Lehner and Döll
710 compiled their database from existing maps, including the GLCC, and inventories. Some
711 wetland types are restricted geographically due to the heterogeneous classification across the
712 source materials. The categories "50-100% wetland" and "25-50% wetland", for example,
713 occur only in North America, "wetland complex" occurs only in Southeast Asia. One
714 consequence is that the global extent of 'bogs, fens, and mires' in the GLWD, 0.8 Mm², is
715 smaller than the Canadian area of peatlands, 1.1 Mm² (Tarnocai et al., 2002), which is
716 dominated by bogs and fens.

Martin Köchy 2015-1-16 10:13

Gelöscht: 5 arc minutes.

Martin Köchy 2015-1-16 10:13

Gelöscht: ..

717 The spatial overlap of the GLWD and the GLCC categories is rather small (Table 6). Only the
718 "Mire bog, fen" category of the GLCC has been adopted completely by the GLWD (Lehner
719 & Döll, 2004). Even categories with similar names like "Freshwater Marsh" vs. "Marsh
720 Wetland" and "Swamp Forest, Flooded Forest" vs. "Wooded Wet Swamps" show little spatial
721 overlap. Despite the GLWD's overall larger wetland area it does not include the areas
722 identified as "rice paddies" in the GLCC.

Martin Köchy 2015-1-16 10:13

Gelöscht: 5

Martin Köchy 2015-1-16 10:13

Gelöscht: category

723 Based on the intersection of GLWD and HWSO (Fig. 3), the global SOC mass in the top 1 m
724 of soil of permanent and non-permanent wetlands (excluding lakes, reservoirs, and rivers) is
725 140 Pg (on 117 Mm² soil area). Using the GLCC Global Ecosystems classification, the area
726 covered by wetlands (excluding inland waters) is much smaller (3 vs 12 Mm²) and contains
727 only 34 Pg SOC (Table 7). The difference is partly due to the classification of large parts of

Martin Köchy 2015-1-16 10:13

Gelöscht: '

Martin Köchy 2015-1-16 10:13

Gelöscht: paddies'

Martin Köchy 2015-1-16 10:13

Gelöscht: 6

735 North America (including the prairie) as temporary or patchy wetland in the GLWD; but even
736 wetlands in a stricter sense cover twice the area and contain nearly twice the mass of SOC in
737 the GLWD compared to the GLCC. Therefore, we combined both maps for the assessment of
738 SOC stocks and masses.

739 Wetlands with the highest C_{org} and highest SOC stocks are bogs, fens, mires, and marshes and
740 the “25–50%” and “50–100%” wetlands in boreal North America. The latter two categories
741 represent mostly bogs, fens, and small lakes. Due to their high C_{org} these wetland types can
742 also be classified as peatland. When wet peatlands are drained, they may no longer qualify as
743 wetlands, but remain peatlands with high C_{org} and a large SOC mass. Drainage exposes the
744 carbon to oxygen and thus accelerates peat decomposition and, depending on circumstances,
745 an increase in BD. The global area of peatland with a minimum peat depth of 30 cm is 3.8

746 Mm^2 based on the International Mire Conservation Group Global Peatland Database (GPD,
747 Joosten, 2010). Total SOC mass of peatlands in the GPD is 447 Pg for their total depth. This
748 estimate is considered conservative because mangroves, salt marshes, paddies, paludified

749 forests, cloud forests, dambos, and Cryosols were omitted because of lack of data. The
750 information available in the database for peatlands is very heterogeneous. For some countries
751 only the total area of peatland is known. When depth information was missing or not
752 plausible, a depth of 2 m was assumed in the GPD, although most peatlands are deeper
753 (Joosten, 2010). It is not clear, which default values were used for C_{org} or BD in the

754 assessment. C_{org} content (organic C fraction of ash-free mass) varies from 0.48–0.52 in
755 *Sphagnum* peat to 0.52–0.59 in *Scheuchzeria* and woody peat (Chambers et al., 2010). Values
756 of BD show much stronger variation. Ash-free bulk density ranged from <0.01 to 0.23 $kg\ dm^{-3}$
757 ³ in 4697 samples (Chambers et al., 2010) with a median of 0.1 $kg\ dm^{-3}$. The variation is due
758 to water content, soil depth, plant material, and degree of decomposition (Boelter, 1968). The
759 highest density is found in well-decomposed, deep peat of herbaceous or woody origin at low

Martin Köchy 2015-1-16 10:13

Gelösch: with

Martin Köchy 2015-1-16 10:13

Gelösch: associated

Martin Köchy 2015-1-16 10:13

Gelösch: cryosols

Martin Köchy 2015-1-16 10:13

Gelösch: concentration (

764 water content. The great variation demands that BD of peatlands actually be measured at
765 several depths and at ambient soil moisture at the same time as the C concentration. If this is
766 not possible, PTFs of BD for peat ought to include water content, decomposition status, and
767 plant material.

768 Peatlands with a certain thickness of organic layer qualify as Histosols. HWSO adopted the
769 FAO definition “Soils having an H horizon of 40 cm or more of organic soil materials (60 cm
770 or more if the organic material consists mainly of sphagnum or moss or has a bulk density of
771 less than 0.1) either extending down from the surface or taken cumulatively within the upper
772 80 cm of the soil; the thickness of the H horizon may be less when it rests on rocks or on
773 fragmental material of which the interstices are filled with organic matter.” (FAO, 1997). The
774 area covered by Histosols in the HWSO (Fig. 4) is 3.3 Mm² (cell area multiplied by fraction
775 of Histosol), slightly lower than the area given by the GPD, and contains 113 Pg SOC. The
776 total area of cells with at least some fraction of Histosol, however, is 10 Mm² containing 188
777 Pg SOC. The area of Histosol outside wetlands (1.7 Mm²) might indicate that a large portion
778 of originally wet peatland has been drained and is exposed to decomposition.

779 The differences in SOC mass estimates between the GLWD and the GLCC indicate that
780 wetland types are defined heterogeneously and that especially the classification of swamp
781 forests, marshes, mangroves, and rice paddies needs to be harmonized. The contrasting land
782 cover classification could be overcome by using the more generic land cover classes
783 developed within the UN Framework Convention on Climate Change (di Gregorio and
784 Jansen, 2005). Remote sensing methods are being developed to improve the mapping of
785 wetlands, e.g., the GlobWetland project (<http://www.globwetland.org>, and Journal of
786 Environmental Management 90, special issue (7)) or the Wetland Map of China (Niu et al.,
787 2009). In situ measurements of soil C_{org}, soil depth, and BD, however, must still be improved,
788 collected, and made available for calculating global SOC mass.

Martin Köchy 2015-1-16 10:13

Gelöscht: .

Martin Köchy 2015-1-16 10:13

Gelöscht: the spatial extent

791 4.4. Carbon in the tropics

792 The high intensity of rain in some parts of the tropics contributes to the presence of wetlands
793 (union of GLWD and GLCC classes as in the previous section) in 9% of the tropical land area
794 (50 Mm² within 23.5°N–23.5°S) containing 40 Pg SOC (Table 8, excluding lakes, reservoirs,
795 rivers). Most of the wetland carbon (27 Pg) is found in marshes and floodplains, and swamp
796 or flooded forests. The GLCC category with the highest SOC mass (10 Pg) is “Rice Paddy
797 and Field” (1.2 Mm² soil and cell area) but only 14% of this area is recognized as wetland in
798 the GLWD.

799 Only 6% of the area of each of the two C-richest tropical wetland types are categorized as
800 Histosols in the HWSD, totaling 0.1 Mm². The total area of Histosol in the HWSD, 0.4 Mm²,
801 agrees with the most recent and detailed, independent estimate of tropical peatland area (Page
802 et al., 2011, defining peatland as soil having >65% organic matter in a minimum thickness of
803 30 cm). The total mass of SOC in grid cells of the spatial layer with at least some fraction of
804 Histosol is 24.2 Pg.

805 Page *et al.* (2011) used peatland area, thickness, BD and C_{org} to calculate the SOC mass for
806 each country within the tropics of Cancer and Capricorn. They tried to trace the original data
807 and used best estimates where data were missing. Most data was available for area, but less
808 data was available for thickness. Page *et al.* (2011) used 25% of maximum thickness when
809 only this information was reported instead of mean thickness and used 0.5 m when no
810 thickness was reported. The percentiles of the frequency distribution of their best estimate of
811 thickness weighted by their best estimate of area per country is 0-10%: 0.5 m, 25%: 1.75 m,
812 50%-90%: 5.5 m, 97.5%: 7.0 m, mean: 4.0 m ± 2.2 m SD. This distribution can be used for
813 estimates of SOC mass and associated uncertainty in other tropical peatlands. Data on BD and
814 SOC concentration were rare. When they were provided they often referred only to the

Martin Köchy 2015-1-16 10:13
Gelöscht: 7

Martin Köchy 2015-1-16 10:13
Gelöscht: pixel

Martin Köchy 2015-1-16 10:13
Gelöscht: C

Martin Köchy 2015-1-16 10:13
Gelöscht: C

819 | subsurface, although these parameters vary with depth. When these data were missing, Page
820 | *et al.* (2011) used 0.09 g cm^{-3} and 56% as best estimates based on literature reviews.

821 | Consequently, their best estimate of SOC mass for tropical peatlands is 88.6 Pg for the whole
822 | soil depth, with a minimum of 81.7 and a maximum of 91.9 Pg. If one assumes an average
823 | peat thickness of 4 m and uniform vertical mass distribution, the top 1 m contains 22 Pg of
824 | SOC, close to our HWSD-based estimate for grid cells containing Histosol (24 Pg). Joosten
825 | (2010) estimated SOC mass for individual tropical countries based on the Global Peatland
826 | Database. For some countries the difference between Joosten's and Page *et al.*'s estimates are
827 | large. For example, Joosten's estimate for Sudan is 1.98 Pg, whereas Page *et al.* have 0.457
828 | Pg. These differences may be caused by different definitions of "peat" and variability in depth
829 | estimates, SOC concentration, and BD in the data sources.

830 | For estimating total tropical SOC mass without depth limit, we add 3/4 of Page *et al.*'s best
831 | estimate for tropical peatland (66.5 Pg) to represent SOC deeper than 1 m to our estimate of
832 | SOC mass in the top 1 m, resulting in 421 Pg. (This addition, however, excludes SOC below
833 | 1 m outside peatlands.) Thus, peatlands contain about 6% of the tropical SOC mass within the
834 | first meter and approximately 21% of the total tropical SOC mass (without depth limit).

835 | Obviously, the uncertainty of these estimates is great.

Martin Köchy 2015-1-16 10:13

Gelöscht: the tropics

Martin Köchy 2015-1-16 10:13

Gelöscht: for the whole soil depth

Martin Köchy 2015-1-16 10:13

Gelöscht: .

839 **5. Conclusions**

840 *5.1. Global carbon mass – reprise*

841 Assuming that the assessment of Tarnocai *et al.* (2009) of the SOC mass in the permafrost
842 region is more accurate than that of HWSD, we update the global SOC mass within the top 1
843 m to 1325 Pg (1062 [HWSD global SOC mass] – 233 [HWSD permafrost [region](#) SOC mass,
844 [Table 4](#)] + 496 [Tarnocai *et al.*'s estimate] Pg). We can use the best estimates of the total
845 SOC mass for the permafrost region (1672 Pg — including deep carbon and high carbon
846 content deposits —, Tarnocai *et al.*, 2009) and the tropics (421 Pg) and add it to the SOC mass
847 outside these areas (473 Pg). This sum (2567 Pg) does not yet comprise SOC below 1 m
848 outside the permafrost region and the tropics (389 Pg, Jobbágy and Jackson, 2000). Thus the
849 total SOC in soil is estimated at about 3000 Pg, but large uncertainties remain, especially for
850 depths >1 m.

851 The BD of peat varies between 0.05 and 0.26 kg/dm³ (Boelter, 1968). If the same range holds
852 for Histosols (3.3 Mm² Histosol area, 1 m depth, 34% C_{org}), this variation alone introduces an
853 uncertainty range of –56 to +180 Pg for the estimate of global SOC in the top meter, which is
854 larger than the estimated annual global soil respiration (79.3–81.8 Pg SOC, Raich *et al.*,
855 2002). The areal extent of peatlands, their depth, and BD should therefore receive the greatest
856 focus of future soil mapping activities.

857 Soil monitoring is crucial for detecting changes in SOC stocks and as a reference for
858 projecting changes in the global carbon pool using models (Wei *et al.*, 2014, Wieder *et al.*,
859 2014, Yan *et al.*, 2014). The following conclusions from our study with respect to improved
860 soil monitoring agree with more comprehensive recommendations by an international group
861 of experts (Jandl *et al.*, 2014). Extra care is necessary to reduce variability of data because

862 variability reduces the potential of detecting change. Classification of soils as it is currently
863 used in mapping produces uncertainty in the reported C stock when the characteristics of soil
864 classes are aggregated and then used in further calculations. The use of pedotransfer rules and
865 functions further increases the uncertainty of the real values. Since pedotransfer functions are
866 entirely empirical in nature, it is preferable that they be derived from soils that are similar in
867 nature to the soils to which the functions will be applied. For purposes of detecting actual
868 change in C stocks their uncertainty should be quantified. Of course it would be best if C_{org},
869 BD, and coarse fragments were measured at the same point or sample to reduce effects of
870 spatial variability. Predictive mapping techniques, including geo-statistics, modelling, and
871 other quantitative methods (McBratney et al., 2003a, [Grunwald et al., 2011](#)), especially in
872 conjunction with proximal (radiometry, NIR spectroscopy) or hyperspectral remote-sensing of
873 soil properties ([Gomez et al., 2008](#), [Stockmann et al., 2015](#)) can potentially reduce
874 uncertainties in SOC mapping introduced by soil classification and help in interpreting spatio-
875 temporal patterns. Whether soils are mapped in the classical way or by predictive methods,
876 mapping of soils should be coordinated with the direct or indirect mapping of SOC input and
877 its controlling factors (land use, land cover, crop type, land use history and land management)
878 and extent and soil depth of wetlands, peatlands, and permafrost.

879 Uncertainty of SOC stocks in current maps could further be reduced if all soil types and
880 regions were well represented by soil profile data with rich soil characteristics. Many soil
881 profile data collected by governments and publicly funded projects remain unused because
882 they are not available digitally; their use is restricted because of data protection issues, or
883 because they are only known to a very limited number of soil scientists. Existing approaches
884 such as the Northern Circumpolar Soil Carbon Base, the GlobalSoilMap.net project, and the
885 Global Soil Partnership (coordinated by FAO), are important steps to improve the situation.

886 These activities would benefit further if all publicly funded, existing soil profile data were
887 made publicly available to the greatest possible extent.

Martin Köchy 2015-1-16 10:13

Gelöscht: were

Martin Köchy 2015-1-16 10:13

Gelöscht: ,

Martin Köchy 2015-1-16 10:13

Gelöscht: be benefitted

891 Another source of uncertainty is introduced because profile data and soil maps have been
892 generated by a multitude of methods. Furthermore, if different methods are preferably used
893 for particular soil types or regions, small differences multiplied by large areas can result in
894 significant differences at the global level. Therefore, international activities to harmonize
895 methods of sampling, calculation, and scaling should be supported. The harmonized methods
896 should then actually be applied in soil sampling. Preferably, samples should be archived so
897 that soils can be reanalyzed with improved or new methods or for checking data by more than
898 one laboratory.

899 5.2. Implications

900 The strong effect of BD values on the calculation of SOC stocks and regional or global
901 masses should guide the focus of global observation networks to improve not only the
902 observation of SOC concentrations but also on BD. Furthermore, our study describes for the
903 first time the frequency distribution of SOC stocks within broad classes of land-use/land-
904 cover and C-rich environments based on one of the most exhaustive, harmonized, spatially
905 explicit global databases available to date. The frequency distribution allows a more focused
906 spatial extrapolation and assessment of accuracy in studies where SOC is used as an
907 independent variable (e.g., Pregitzer and Euskirchen, 2004). The frequency distributions also
908 provide a foundation for targeting SOC conservation measures (Powlson et al., 2011) and for
909 improving carbon accounting methods with associated uncertainties as used in the UNFCCC
910 (García-Oliva and Masera, 2004).
911 CO₂ emissions from soils are used in calculations of the global carbon cycle. Direct
912 observations of CO₂ emissions from soils (e.g., by eddy-flux towers), however, cannot be
913 implemented in a spatially contiguous way. Indirect measurements by remote sensing can
914 improve the spatial coverage but require ground observations for conversion from observed

Martin Köchy 2015-1-16 10:13

Gelösch: Our

Martin Köchy 2015-1-16 10:13

Gelösch: .

Martin Köchy 2015-1-16 10:13

Gelösch: classes

Martin Köchy 2015-1-16 10:13

Gelösch: The strong effect of BD values on SOC stocks and regional or global masses guides the focus of global observation networks to improve not only the observation of SOC concentrations but also on BD.

923 radiation to loss of CO₂ from soils and distinction from other CO₂ sources (Ciais et al., 2010).
924 At the global scale, in-situ measurements must be complemented by modelling activities,
925 which are greatly improved if variation in key factors like SOC can be accounted for. Thus,
926 more detailed information on the global distribution of SOC, horizontally and vertically,
927 including accounts of its accuracy and its variability, are necessary to improve estimates of
928 the global carbon flow.

Martin Köchy 2015-1-16 10:13

Gelösch: soil organic carbon

Martin Köchy 2015-1-16 10:13

Gelösch: can

929 **Author contributions**

930 MK designed and carried out the analyses and wrote the manuscript, RH contributed a
931 thorough analysis of inconsistencies in the HWSD and alternative estimates, AF suggested the
932 topic and provided valuable insights on the presentation of the data.

933 Acknowledgments. We thank Charles Tarnocai for comments on the manuscript. We
934 appreciate the constructive comments of the reviewers. MK was funded by EU FP7 project
935 COCOS (grant 212196) and FACCE MACSUR (BMBF grant 031A103A).

938 **References**

- 939 Amundson, R.: The carbon budget in soils, *Annual Review of Earth and Planetary Sciences*, 29, 535-
940 562, doi:10.1146/annurev.earth.29.1.535, 2001.
- 941 Batjes, N. H.: Total carbon and nitrogen in the soils of the world, *European Journal of Soil Science*,
942 47, 151-163, doi:10.1111/j.1365-2389.1996.tb01386.x, 1996.
- 943 Batjes, N. H.: Harmonized soil profile data for applications at global and continental scales: updates to
944 the WISE database, *Soil Use and Management*, 25, 124-127, doi:10.1111/j.1475-
945 2743.2009.00202.x, 2009.
- 946 Boelter, D. H.: Important physical properties of peat materials, *Proceedings of the Third International*
947 *Peat Congress*, Ottawa, Ontario, Canada, 18–23 August 1968, Canada. Dept. of Energy, Mines
948 and Resources and National Research Council of Canada, 1968.
- 949 Carré, F., R. Hiederer, V. Blujdea, and R. Koeble: Background guide for the calculation of land
950 carbon stocks in the biofuels sustainability scheme drawing on the 2006 IPCC Guidelines for
951 National Greenhouse Gas Inventories, JRC Scientific and Technical Reports, EUR 24573 EN,
952 Office for Official Publications of the European Communities, Luxembourg, Office for Official
953 Publications of the European Communities, available at:
954 http://eusoils.jrc.ec.europa.eu/ESDB_Archive/eusoils_docs/other/EUR24573.pdf, 2010.
- 955 Chambers, F. M., Beilman, D. W., and Yu, Z.: Methods for determining peat humification and for
956 quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate
957 and peatland carbon dynamics, *Mires and Peat*, 7, 7.1-7.10, 2010/2011.
- 958 Ciais, P., A. J. Dolman, R. Dargaville, L. Barrie, A. Bombelli, J. Butler, P. Canadell, and T.
959 Moriyama: GEO Carbon Strategy, GEO Secretariat and FAO, Geneva and Rome, GEO
960 Secretariat and FAO, available at:
961 http://www.earthobservations.org/documents/sbas/cl/201006_geo_carbon_strategy_report.pdf,
962 2010.

963 Ellert, B. H., H. H. Janzen, and B. G. McConkey: Measuring and comparing soil carbon storage, in:
964 Assessment methods for soil carbon, *Advances in Soil Science*, Lal, R., J. M. Follett, and B. A.
965 Stewart (Eds), Lewis, Boca Raton, Florida, 131-146, 2001.

966 FAO: FAO/Unesco Soil Map of the World, Revised Legend, with corrections and updates. Originally
967 published in 1988 as *World Soil Resources Report 60*, FAO, Rome. Reprinted with updates,
968 Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at:
969 http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997.

970 Digital Soil Map of the World, Rome, Italy, available at:
971 <http://www.fao.org/geonetwork/srv/en/metadata.show?id=14116>, 2007.

972 FAO, IIASA, ISRIC, ISSCAS, and JRC: [Harmonized World Soil Database \(version 1.1\)](#), FAO and
973 IIASA, Rome, Italy, and Laxenburg, Austria, 2009.

974 [FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database \(version 1.2\), FAO and](#)
975 [IIASA, Rome, Italy, and Laxenburg, Austria, 2012.](#)

976 Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuisen, L. Verelst, and D. Wiberg: Global agro-
977 ecological zones assessment for agriculture (GAEZ 2008), IIASA, Laxenburg, Austria and
978 FAO, Rome, Italy., IIASA, Laxenburg, Austria and FAO, Rome, Italy., available at:
979 <http://www.iiasa.ac.at/Research/LUC/External-World-soil->
980 [database/HTML/SoilQualityData.html?sb=11](http://www.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/SoilQualityData.html?sb=11), 2008.

981 Freeman, C., Ostle, N., and Kang, H.: An enzymic 'latch' on a global carbon store, *Nature*, 409, 149,
982 doi:10.1038/35051650, 2001.

983 García-Oliva, F., and Masera, O. R.: Assessment and Measurement Issues Related to Soil Carbon
984 Sequestration in Land-Use, Land-Use Change, and Forestry (LULUCF) Projects under the
985 Kyoto Protocol, *Clim. Change*, 65, 347-364, doi:10.1023/B:CLIM.0000038211.84327.d9, 2004.

986 Global Soil Data Task Group: Global Soil Data Products CD-ROM (IGBP-DIS), Available from Oak
987 Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
988 [<http://www.daac.ornl.gov>], 2000.

Martin Köchy 2015-1-16 10:13

Gelöscht: .

990 [Gomez, C., Viscarra, R., Raphael A., and McBratney, A. B.: Soil organic carbon prediction by](#)
991 [hyperspectral remote sensing and field vis-NIR spectroscopy: An Australian case study,](#)
992 [Geoderma, 146, 403-411, doi:10.1016/j.geoderma.2008.06.011, 2008.](#)

993 Gorham, E.: Northern peatlands: role in the carbon cycle and probable responses to climatic warming,
994 *Ecol. Appl.*, 1, 182-195, doi:10.2307/1941811, 1991.

995 GRASS Development Team: Geographic Resources Analysis Support System (GRASS) Software,
996 Version 6.4.2, Open Source Geospatial Foundation, <http://grass.osgeo.org>, 2011.

997 di Gregorio, A., and L. J. M. Jansen: Land cover classification system (LCCS). Classification concepts
998 and user manual. Software version (2), Food and Agriculture Organization, Rome, Italy, Food
999 and Agriculture Organization, available at:
1000 <http://www.fao.org/docrep/008/y7220e/y7220e00.htm#Contents>, 2005.

1001 [Grunwald, S., Thompson, J. A., and Boettinger, J. L.: Digital soil mapping and modeling at](#)
1002 [continental scales: finding solutions for global issues, Soil Sci. Soc. Am. J., 75, 1201-1213,](#)
1003 [doi:10.2136/sssaj2011.0025, 2011.](#)

1004 Heginbottom, J. A., J. F. Brown, E. S. Melnikov, and O. J. Ferrians: Circum-arctic map of permafrost
1005 and ground ice conditions, Proceedings of the Sixth International Conference on Permafrost,
1006 Wushan, Guangzhou, China, 2, 1132-1136, South China University Press, 1993.

1007 Henry, M., Valentini, R., and Bernoux, M.: Soil carbon stocks in ecoregions of Africa, *Biogeosciences*
1008 *Discussions*, 6, 797-823, doi:10.5194/bgd-6-797-2009, 2009.

1009 [Hiederer, R.: Data update and model revision for soil profile analytical database of Europe of](#)
1010 [measured parameters \(SPADE/M2\), JRC Scientific and Technical Reports, EUR 24333 EN,](#)
1011 [Office for Official Publications of the European Communities, Luxembourg, Office for Official](#)
1012 [Publications of the European Communities, doi:10.2788/85262, 2010.](#)

1013 [Hiederer, R.](#), and M. Köchy: Global soil organic carbon estimates and the Harmonized World Soil
1014 Database, JRC Scientific and Technical Reports, 68528/EUR 25225 EN, Joint Research Centre,
1015 Ispra, Italy, Joint Research Centre, doi:10.2788/13267, 2011.

1016 Hiederer, R., F. Ramos, C. Capitani, R. Koeble, V. Blujdea, O. Gomez, D. Mulligan, and L. Marelli:
1017 Biofuels: a new methodology to estimate GHG emissions from global land use change, JRC

Martin Köchy 2015-1-16 10:13

Gelöscht: /05/01

1019 Scientific and Technical Reports, EUR 24483 EN, Office for Official Publications of the
1020 European Communities, Luxembourg, Office for Official Publications of the European
1021 Communities, [doi:10.2788/48910](https://doi.org/10.2788/48910), 2010.

1022 IUSS Working Group WRB: World reference base for soil resources 2006. A framework for
1023 international classification, correlation and communication, World Soil Resources Reports, 103,
1024 Food and Agriculture Organization of the United Nations, Rome, 2006.

1025 Jandl, R., Rodeghiero, M., Martinez, C., Cotrufo, M. F., Bampa, F., van Wesemael, B., Harrison, R.,
1026 B., Guerrini, I. A., Richter, D. d., Jr., Rustad, L., Lorenz, K., Chabbi, A., and Miglietta, F.:
1027 Current status, uncertainty and future needs in soil organic carbon monitoring, Science of the
1028 Total Environment, 468, 376-383, doi:10.1016/j.scitotenv.2013.08.026, 2014.

1029 Jobbágy, E. G., and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to
1030 climate and vegetation, Ecol. Appl., 10, 423-436, doi:10.1890/1051-
1031 0761(2000)010[0423:TVDOS0]2.0.CO;2, 2000.

1032 Joosten, H.: The global peatland CO₂ picture. Peatland status and emissions in all countries of the
1033 world, Wetlands International, Ede, 2010.

1034 Köchy, M., and A. Freibauer: Global spatial distribution of wetlands, COCOS Report, D4.3a, Johann
1035 Heinrich von Thünen-Institut, Braunschweig, Germany, Johann Heinrich von Thünen-Institut,
1036 available at: http://www.cocos-carbon.org/docs/D4.3a_wetlands_report.pdf, 2009.

1037 Lehner, B., and Döll, P.: Development and validation of a global database of lakes, reservoirs and
1038 wetlands, J. Hydrol., 296, 1-22, doi:10.1016/j.jhydrol.2004.03.028, 2004.

1039 Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, J., Yang, L., and Merchant, J. W.:
1040 Development of a Global Land Cover Characteristics Database and IGBP DISCover from 1-km
1041 AVHRR Data, International Journal of Remote Sensing, 21, 1303-1330,
1042 doi:10.1080/014311600210191, 2000.

1043 Mäkilä, M.: Calculation of the energy content of mires on the basis of peat properties [In Finnish with
1044 English summary], Report of Investigation, 121, Geological Survey of Finland, Geological
1045 Survey of Finland, 1994.

1046 McBratney, A. B., Mendonça, S., M.L., and Minasny, B.: On digital soil mapping, *Geoderma*, 117, 3-
1047 52, doi:10.1016/S0016-7061(03)00223-4, 2003/11//a.

1048 McBratney, A. B., Mendonça Santos, M. L., and Minasny, B.: On digital soil mapping, *Geoderma*,
1049 117, 3-52, doi:10.1016/S0016-7061(03)00223-4, 2003b.

1050 Mitra, S., Wassmann, R., and Vlek, P.: An appraisal of global wetland area and its organic carbon
1051 stock, *Current Science*, 88, 25-35, 2005.

1052 Nachtergaele, F. O.: From the Soil Map of the World to the Digital Global Soil and Terrain Database:
1053 1960–2002, in: *Handbook of Soil Science*, Sumner, M. E. (Ed.), CRC Press, Boca Raton, H5-
1054 17, 1999.

1055 Niu, Z. G., Gong, P., Cheng, X., Guo, J. H., Wang, L., Huang, H. B., Shen, S. Q., Wu, Y. Z., Wang,
1056 X. F., Wang, X. W., Ying, Q., Liang, L., Zhang, L. N., Wang, L., Yao, Q., Yang, Z. Z., Guo, Z.
1057 Q., and Dai, Y. J.: Geographical analysis of China's wetlands preliminarily derived from
1058 remotely sensed data, *Science in China – Series D: Earth Sciences*, 39, 188-203, 2009.

1059 Page, S. E., Rieley, J. O., and Banks, C. J.: Global and regional importance of the tropical peatland
1060 carbon pool, *Glob. Change Biol.*, 17, 798-818, doi:10.1111/j.1365-2486.2010.02279.x, 2011.

1061 Powlson, D. S., Whitmore, A. P., and Goulding, K. W. T.: Soil carbon sequestration to mitigate
1062 climate change: a critical re-examination to identify the true and the false, *European Journal of*
1063 *Soil Science*, 62, 42-55, doi:10.1111/j.1365-2389.2010.01342.x, 2011.

1064 Pregitzer, K. S., and Euskirchen, E. S.: Carbon cycling and storage in world forests: biome patterns
1065 related to forest age, *Glob. Change Biol.*, 10, 2052-2077, doi:10.1111/j.1365-
1066 2486.2004.00866.x, 2004.

1067 R Development Core Team: *R: A language and environment for statistical computing*, R Foundation
1068 for Statistical Computing, Vienna, 2011.

1069 Raich, J. W., Potter, C. S., and Bhagawati, D.: Interannual variability in global soil respiration, 1980–
1070 94, *Glob. Change Biol.*, 8, 800-812, doi:10.1046/j.1365-2486.2002.00511.x, 2002.

1071 Reich, P.: Soil organic carbon map, USDA-NRCS,
1072 <http://soils.usda.gov/use/worldsoils/mapindex/soc.html>, last access: 2011-09-20, 2000.

1073 Scharlemann, J. P. W., Tanner, E., V. J., Hiederer, R., and Kapos, V.: Global soil carbon:
1074 understanding and managing the largest terrestrial carbon pool, *Carbon Management*, 5, 81-91,
1075 doi:10.4155/CMT.13.77, 2014.

1076 Shangguan, W., Dai, Y., Duan, Q., Liu, B., and Yuan, H.: A global soil data set for earth system
1077 modeling, *Journal of Advances in Modeling Earth Systems*, 6, 249-263,
1078 doi:10.1002/2013MS000293, 2014.

1079 [Stockmann, U., Malone, B. P., McBratney, A. B., and Minasny, B.: Landscape-scale exploratory](#)
1080 [radiometric mapping using proximal soil sensing, *Geoderma*, 239–240, 115-129,](#)
1081 [doi:10.1016/j.geoderma.2014.10.005, 2015.](#)

1082 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic
1083 carbon pools in the northern circumpolar permafrost region, *Glob. Biogeochem. Cycles*, 23,
1084 GB2023, doi:10.1029/2008GB003327, 2009.

1085 Tarnocai, C., I. M. Kettles, and B. Lacelle: Peatlands of Canada database, Open File, 4002, Geological
1086 Survey of Canada, doi:10.4095/213529, 2002.

1087 Turunen, J.: Development of Finnish peatland area and carbon storage 1950–2000, *Boreal*
1088 *Environment Research*, 13, 319-334, 2008.


1089 Wei, X., Shao, M., Gale, W., and Li, L.: Global pattern of soil carbon losses due to the conversion of
1090 forests to agricultural land, *Scientific Reports*, 4, 4062, 2014.

1091 Wieder, W. R., Boehnert, J., and Bonan, G. B.: Evaluating soil biogeochemistry parameterizations in
1092 Earth system models with observations, *Glob. Biogeochem. Cycles*, 28, 211-222,
1093 doi:10.1002/2013GB004665, 2014.

1094 Yan, Y., Luo, Y., Zhou, X., and Chen, J.: Sources of variation in simulated ecosystem carbon storage
1095 capacity from the 5th Climate Model Intercomparison Project (CMIP5), *Tellus Series B –*
1096 *Chemical and Physical Meteorology*, 66, 22568, doi:10.3402/tellusb.v66.22568, 2014.

1097 Zimov, S. A., Davydov, S. P., Zimova, G. M., Davydova, A. I., Schuur, E. A. G., Dutta, K., and
1098 Chapin, F. S.: Permafrost carbon: Stock and decomposability of a globally significant carbon
1099 pool, *Geophys. Res. Lett.*, 33, L20502, doi:10.1029/2006GL027484, 2006.

1100



1102 Table 1. Definition of terms with respect to organic soil carbon.

| <u>Term</u> | <u>Definition</u> |
|-------------------------------------|---|
| <u>Concentration</u> | organic carbon mass/soil dry mass, C_{org} |
| <u>Areal density (of fine soil)</u> | $C_{org} \times \text{depth} \times (1 - \text{fractional volume of rocks, coarse roots, and ice})$ |
| <u>Stock</u> | areal density of fine soil integrated over all layers to a specified depth |
| <u>Mass</u> | stock integrated over a specified area |

1104

1105

Martin Köchy 2015-1-16 10:13
Gelöscht: Term - Definition - ... [10]

1108 **Table 2.** Changes to the global SOC mass in the top 1 m after modifications to the HWSD 1.1
 1109 database.

| processing step | SOC mass (Pg) |
|---|---------------|
| no modification | 2469.5 |
| (1) filling of missing values for C _{org} | 2470.6 |
| (2) filling of missing values for BD | 2471.3 |
| (3) adjusting BD values when C _{org} >3% | 1230.2 |
| (-) replacing BD values only for Histosols | 1113.3 |
| (4) adjusting BD values for C _{org} >3% & replacing BD for Histosols | 1060.9 |

1111

1112
 1113 **Table 3.** Soil organic carbon stocks by continent. For the definition of ‘continents’ we used
 1114 the ESRI (2002) map of continents with coastlines extended by 2 pixels to increase the
 1115 overlap. $1 \text{ Mm}^2 = 10^6 \text{ km}^2$

| Continent converted to 30" raster | Soil area (Mm²) | Carbon stock, 0–1 m (Pg) HWSD v.1.1-modified |
|--|-----------------------------------|--|
| Asia, incl. Malay Archipelago | 42.0 | 369 |
| North America, incl. Greenland, Central America | 21.3 | 223 |
| Europe, incl. Iceland, Svalbard, Novaya Zemlya | 9.4 | 110 |
| Africa, incl. Madagascar | 27.2 | 148 |
| South America | 17.7 | 163 |
| Australia, New Zealand, Pacific Islands | 8.0 | 46 |
| non-overlapping pixels | 0.2 | 2 |
| total (90°N – 60°S) | 125.8 | 1061 |

1116

1117 Table 4 Organic carbon mass (top 1 m) of soils with gelic properties in HWSD v.1.1-
 1118 modified. (All areas north of 60°S). Percentiles refer to the distribution of C stocks in each
 1119 cell within the soil area mentioned. 1 Mm² = 10⁶ km². Hist/soil: fraction of soil area covered
 1120 by Histosols.

| Gelic phase | Cell area (Mm ²) | Soil area (Mm ²) | Hist/soil | C stock (kg m ⁻²), percentiles | | | | | C mass (Pg) |
|-----------------------|------------------------------|------------------------------|-----------|--|------|------|------|------|-------------|
| | | | | 5% | 25% | 50% | 75% | 95% | |
| | | | | continuous, >90% of area | 5.46 | 5.30 | 12% | 5.9 | |
| discontinuous, 50–90% | 4.11 | 4.07 | 12% | 6.4 | 6.5 | 9.5 | 15.8 | 28.9 | 51.8 |
| sporadic, 10–50% | 3.79 | 3.68 | 6% | 3.8 | 8.3 | 12.5 | 15.6 | 19 | 45.3 |
| isolated, 0–10% | 0.05 | 0.05 | 86% | 8.4 | 27.9 | 32.8 | 32.8 | 32.8 | 1.5 |
| whole area | 13.41 | 13.10 | 11% | 5.3 | 6.9 | 9.8 | 15.6 | 30.6 | 163.8 |

1121

Martin Köchy 2015-1-16 10:13
 Gelösch: 3

Martin Köchy 2015-1-16 10:13
 Gelösch: pixel

Martin Köchy 2015-1-16 10:13
 Gelösch: gelic

Martin Köchy 2015-1-16 10:13
 Gelösch: pixel

Martin Köchy 2015-1-16 10:13
 Gelösch: soil

Martin Köchy 2015-1-16 10:13
 Gelösch: hist

1128 | Table 5. Comparison of organic carbon stocks (top 1 m) between HWSD v.1.1-modified and
 1129 | NCSCDB (Tarnocai *et al.* 2009). Permafrost contingency refers to the Circumarctic
 1130 | Permafrost Map. NCSCDB used different soil areas than HWSD. Percentiles refer to the
 1131 | distribution of C stocks in each **grid cell** within the soil area mentioned. 1 Mm² = 10⁶ km².

| Permafrost contingency of NCSCDB | HWSD | | | | | NCSCDB | | | | | | |
|----------------------------------|------------------------------|------------------------------|--|-----|------|--------|------|-------------|------------------------------|-------------------------------------|-------------|--|
| | Cell area (Mm ²) | Soil area (Mm ²) | C stock (kg m ⁻²), percentiles | | | | | C mass (Pg) | Soil area (Mm ²) | C stock (kg m ⁻²), mean | C mass (Pg) | |
| | | | 5% | 25% | 50% | 75% | 95% | | | | | |
| continuous, >90% of area | 10.64 | 9.97 | 4.1 | 6.5 | 8 | 14.6 | 18.8 | 105.8 | 10.1 | 29.5 | 299 | |
| discontinuous, 50–90% | 3.17 | 3.05 | 4.4 | 6.9 | 12.9 | 16.9 | 32.6 | 41.3 | 3.1 | 21.8 | 67 | |
| sporadic, 10–50% | 3.08 | 2.94 | 4.9 | 7.4 | 12.7 | 17 | 35.5 | 40.3 | 2.6 | 24.3 | 63 | |
| isolated, 0–10% | 3.67 | 3.55 | 5.6 | 7.8 | 10.1 | 16 | 32.3 | 45.4 | 3.0 | 22.6 | 67 | |
| whole area | 20.55 | 19.52 | 4.4 | 6.9 | 9.4 | 15.5 | 28 | 232.7 | 18.8 | 26.4 | 496 | |

1132

Martin Köchy 2015-1-16 10:13
 Gelösch: 4

Martin Köchy 2015-1-16 10:13
 Gelösch: pixel

Martin Köchy 2015-1-16 10:13
 Gelösch: permafrost

Martin Köchy 2015-1-16 10:13
 Gelösch: pixel

Martin Köchy 2015-1-16 10:13
 Gelösch: soil

Martin Köchy 2015-1-16 10:13
 Gelösch: soil

1139 Table 6. Area and spatial overlap of wetland types in GLWD and GLCC (grid cell area, Mm²)
 1140 within the extent of the HWSD.

| GLWD | GLCC, ecosystems legend | | | | | | | |
|------------------------------------|-------------------------|-----------------|------------------|----------------------|--------------|-------------------|-------------------------|----------|
| | area (Mm ²) | 14 Inland Water | 45 Marsh Wetland | 13 Wooded Wet Swamps | 72 Man-grove | 44 Mire, Bog, Fen | 36 Rice Paddy and Field | Dry-land |
| 1-3 Lake, Reservoir, River | <u>2.370</u> | 1.437 | 0.000 | 0.002 | 0.006 | 0.027 | 0.008 | 0.845 |
| 4 Freshwater Marsh, Floodplain | <u>2.487</u> | 0.077 | 0.015 | 0.003 | 0.006 | 0.058 | 0.167 | 2.155 |
| 5 Swamp Forest, Flooded Forest | <u>1.154</u> | 0.041 | - | 0.013 | 0.001 | - | 0.006 | 1.090 |
| 6 Coastal Wetland | <u>0.413</u> | 0.015 | 0.001 | 0.007 | 0.011 | 0.002 | 0.026 | 0.321 |
| 7 Pan, Brackish/Saline Wetland | <u>0.433</u> | 0.002 | <0.001 | <0.001 | <0.001 | - | 0.001 | 0.429 |
| 8 Bog, Fen, Mire | <u>0.710</u> | - | - | - | - | 0.710 | - | - |
| 9 Intermittent Wetland/Lake | <u>0.689</u> | 0.004 | <0.001 | <0.001 | <0.001 | - | 0.003 | 0.681 |
| 10 50-100% Wetland | <u>1.762</u> | 0.045 | - | 0.005 | - | - | - | 1.693 |
| 11 25-50% Wetland | <u>3.153</u> | 0.065 | - | <0.001 | - | - | - | 3.077 |
| 12 Wetland Complex (0-25% Wetland) | <u>0.898</u> | <0.001 | - | - | - | - | 0.046 | 0.846 |
| Dryland | <u>120.433</u> | 0.646 | 0.045 | 0.052 | 0.024 | - | 2.149 | 116.896 |

Martin Köchy 2015-1-16 10:13
Gelöscht: 5

Martin Köchy 2015-1-16 10:13
Gelöscht: Spatial

Martin Köchy 2015-1-16 10:13
Gelöscht: .

Martin Köchy 2015-1-16 10:13
Eingefügte Zellen

Martin Köchy 2015-1-16 10:13
Gelöscht: Dryland

Martin Köchy 2015-1-16 10:13
Gelöscht: -

Martin Köchy 2015-1-16 10:13
Eingefügte Zellen

Martin Köchy 2015-1-16 10:13
Eingefügte Zellen

1141

1142

1148
1149
1150
1151
1152
1153
1154
1155

Table 7. Organic carbon stocks and masses in the top 1 m of *global* wetland soils derived from the HWSO v1.1-modified. Wetland extent is primarily according to the Global Lake and Wetlands Database (1–12), augmented by wetland in the GLCC (13–72). Percentiles refer to the distribution of C stocks in each *grid cell* within the soil area mentioned. **C mass of permanent wetlands (types B–D) is 81.8 Pg, that of all wetlands except open waters (types B–K) is 158.1 Pg.** 1 Mm² = 10⁶ km². *Hist/soil*: fraction of soil area covered by Histosols.

| Wetland type | Cell area | Soil area | Hist. / soil | C stock (kg m ⁻²), percentiles | | | | | C mass (Pg) |
|---|--------------------|--------------------|--------------|--|------|------|------|------|-------------|
| GLWD and GLCC category | (Mm ²) | (Mm ²) | % | 5 % | 25 % | 50 % | 75 % | 95 % | |
| A 1–3 Lake, Reservoir, River 14 Inland Water | 3.01 | 2.11 | 7 | 4.2 | 6.5 | 9 | 14 | 24 | 22 |
| B 4 Freshwater Marsh, Floodplain 45 Marsh Wetland | 2.53 | 2.48 | 17 | 4.4 | 7 | 10 | 19 | 38 | 32 |
| 5 Swamp Forest, Flooded | 1.21 | 1.21 | 6 | 3.6 | 5.6 | 8.6 | 13 | 33 | 13 |
| C Forest 13 Wooded Wet Swamps | | | | 6 | | | 6 | 8 | 2 |
| D 8/44 Bog, Fen, Mire | 0.71 | 0.68 | 14 | 4.4 | 8.4 | 14 | 18 | 35 | 10 |
| E 7 Pan, Brackish/ Saline Wetland | 0.43 | 0.31 | <1 | 2.8 | 4 | 4.7 | 5.4 | 7.5 | 1.5 |
| F 6 Coastal Wetland 72 Mangrove | 0.44 | 0.43 | 4 | 3.9 | 6.1 | 7.3 | 11 | 21 | 4.4 |
| G 36 Rice Paddy and Field | 2.15 | 2.14 | <1 | 4.7 | 6 | 7.1 | 8.9 | 12 | 17 |
| H 9 Intermittent Wetland/Lake | 0.69 | 0.60 | <1 | 2.3 | 3.6 | 4.4 | 5.9 | 9.6 | 3 |
| I 10 50-100% Wetland | 1.75 | 1.74 | 33 | 6.9 | 12 | 13 | 24 | 38 | 31 |
| J 11 25-50% Wetland | 3.14 | 3.11 | 10 | 5.6 | 8.8 | 12 | 14 | 28 | 38 |
| K 12 Wetland Complex (0- 25% Wetland) | 0.9 | 0.89 | 1 | 5.8 | 5.9 | 5.9 | 7.3 | 12 | 6.7 |
| Dryland | 117.2 | 110.1 | 2 | 2.4 | 4.9 | 7.1 | 10 | 18 | 880 |
| | 4 | 5 | 5 | | | | 3 | 1 | |

Martin Köchy 2015-1-16 10:13
Gelöscht: 6

Martin Köchy 2015-1-16 10:13
Gelöscht: pixel

Martin Köchy 2015-1-16 10:13
Gelöscht: Pixel

Martin Köchy 2015-1-16 10:13
Gelöscht: 2

Martin Köchy 2015-1-16 10:13
Gelöschte Zellen

Martin Köchy 2015-1-16 10:13
Gelöschte Zellen

Martin Köchy 2015-1-16 10:13
Gelöscht: 81.8

Martin Köchy 2015-1-16 10:13
Gelöscht: 158.1

1156

1157

1164

1165 Table 8. Organic carbon stocks and masses in the top 1 m of tropical wetland soils derived

1166 from HWSD v.1.1-modified. Wetlands classified primarily according to the Global Lake and

1167 Wetlands Database (1–12), augmented by wetland classes in the GLCC (13–72). Percentiles

1168 refer to the distribution of C stocks in each grid cell within the soil area mentioned. C mass of

1169 permanent wetlands (types B–H) is 38.3 Pg, that of all wetlands except open waters (types

1170 B–K) is 39.9 Pg. 1 Mm² = 10⁶ km². Hist/soil: fraction of soil area covered by Histosols.

Martin Köchy 2015-1-16 10:13
Gelöscht: 7

Martin Köchy 2015-1-16 10:13
Gelöscht: pixel

Martin Köchy 2015-1-16 10:13
Gelöscht: 1 Mm² = 10⁶ km².

| Wetland type | Cell area | Soil area | Hist t./soil | C stock (kg m ⁻²), percentiles | | | | | C mass (Pg) |
|---|--------------------|--------------------|--------------|--|-----|-----|------|------|-------------|
| GLWD and GLCC category | (Mm ²) | (Mm ²) | % | 5% | 25% | 50% | 75% | 95% | |
| A 1–3 Lake, Reservoir, River 14 Inland Water | 0.76 | 0.49 | 2% | 3.9 | 5.9 | 7.9 | 10.6 | 18.8 | 4.5 |
| B 4 Freshwater Marsh, Floodplain | 1.27 | 1.26 | 6% | 3.7 | 6.2 | 7.7 | 10.3 | 24.2 | 12.0 |
| 45 Marsh Wetland | | | | | | | | | |
| C 5 Swamp Forest, Flooded Forest | 1.21 | 1.20 | 6% | 3.6 | 5.6 | 8.6 | 13.6 | 33.8 | 13.2 |
| 13 Wooded Wet Swamps | | | | | | | | | |
| D 8/44 Bog, Fen, Mire | 0.0 | 0.00 | 0% | 2.5 | 6.0 | 6.0 | 11.9 | 12.0 | 0.0 |
| E 7 Pan, Brackish/ Saline Wetland | 0.12 | 0.10 | 0% | 2.5 | 3.2 | 4.3 | 5.3 | 7.5 | 0.5 |
| F 6 Coastal Wetland 72 Mangrove | 0.31 | 0.31 | 4% | 4.0 | 6.1 | 8.5 | 13.7 | 25.7 | 3.4 |
| G 36 Rice Paddy and Field | 1.06 | 1.06 | 1% | 5.1 | 6.2 | 6.9 | 8.1 | 13.2 | 8.4 |
| H 9 Intermittent Wetland/Lake | 0.22 | 0.20 | 0% | 2.2 | 3.3 | 4.1 | 5.0 | 6.4 | 0.8 |
| K 12 Wetland Complex (0- 25% Wetland) | 0.2 | 0.20 | 3% | 5.0 | 5.9 | 6.5 | 8.2 | 13.2 | 1.6 |
| Dryland | 44.71 | 43.06 | 1% | 2.2 | 4.3 | 6.1 | 8.5 | 15.2 | 310.6 |
| Tropical area | 49.87 | 47.88 | 1% | | | | | | 354.9 |

Martin Köchy 2015-1-16 10:13
Gelöscht: Pixel

Martin Köchy 2015-1-16 10:13
Gelöschte Zellen

Martin Köchy 2015-1-16 10:13
Gelöschte Zellen

Martin Köchy 2015-1-16 10:13
Gelöscht: 38.3

Martin Köchy 2015-1-16 10:13
Gelöscht: 39.9

1177 **Figure captions**

1178 | Figure 1. Global stock (a) and mass (b, per 5° latitude) of organic carbon in the top 1 m of the
1179 terrestrial soil calculated from HWSD v.1.1-modified.

Martin Köchy 2015-1-16 10:13
Gelöscht: 100 cm

1180

1181 | Figure 2. Extent of permafrost in HWSD v.1.1. Colour scale: fraction of soil units within a
1182 | 30" grid cell with 'gelic phase' (averaged for display to 30' resolution); red outline:
1183 permafrost attribute in HWSD supplementary data sets SQ1-7 at 5' resolution.

Martin Köchy 2015-1-16 10:13
Gelöscht: pixel

1184

1185 | Figure 3. (a) Global distribution of important wetlands (by carbon mass) according to the
1186 Global Lakes and Wetlands Database and Global Land Cover Characterization. The most
1187 frequent wetland type is displayed within a 0.5° grid cell. Wetland types A-K are explained in
1188 Table 6. (b) Carbon mass in wetland soils (top 1 m) in bands of 5° latitude (calculated from
1189 HWSD v.1.1-modified). (c) Carbon mass in aggregated types of wetland soils (panel b).

1190

1191 | Figure 4. Fraction of Histosol area per 0.5° grid cell according to HWSD v.1.1.

A good knowledge of the global SOC mass and its spatial distribution is necessary for assessing in an international context which soils are most vulnerable or might provide the best opportunity for C sequestration in mitigation of rising greenhouse gas concentrations. At the global scale, in-situ measurements must be complemented by modelling activities, which are greatly improved if variation in key factors like soil organic carbon can be accounted for. In this paper we review existing spatial estimates of SOC stocks and masses, including their uncertainties and underlying methods for estimating the stocks. Our paper reports for the first time area-weighted frequency distributions of carbon stocks within land-use and land-cover classes, using best estimates from several sources. We focus on the large SOC stocks in wetlands, tropical soil, and permafrost at high latitudes and present frequency distributions of SOC stocks within classes of land use, land cover, and geographic region. Furthermore, we provide recommendations for improving global soil mapping.

Methods

Characterization of the Harmonized Word Soil Database

Our analysis of SOC stocks and masses is based on the

Results and Discussion

Global carbon mass

Historic estimates of global SOC mass represented by 27 studies range between 504 and 3000 Pg with a median of 1461 Pg

1.

Seite 13: [3] Auf Seite 10 verschoben (Verschiebung Nr. 10)
2015 · 01 · 16 10:13

Martin Köchy

(Scharlemann et al., 2014).

Seite 13: [3] Auf Seite 10 verschoben (Verschiebung Nr. 10)
2015 · 01 · 16 10:13

Martin Köchy

(Scharlemann et al., 2014).

Seite 13: [4] Gelöscht

Martin Köchy

2015 · 01 · 16 10:13

Before the publication of the HWSD, many global estimates were based on the [Digital] Soil Map of the World ([D]SMW) (FAO, 1997, FAO, 2007). Batjes (1996), using information from 4353 WISE profiles, reported a range of 1462–1548 Pg for 0–1 m depth and 2376–2456 Pg for 1–2 m depth.

1.1.

Seite 13: [5] Auf Seite 9 verschoben (Verschiebung Nr. 7)
2015 · 01 · 16 10:13

Martin Köchy

The International Geosphere Biosphere Program (IGBP) (Global Soil Data Task Group, 2000) produced a map of SOC stock on a 5' by 5' grid derived from the DSMW in conjunction with WISE data (v.1, 1125 profiles).

1.2.

Seite 13: [5] Auf Seite 9 verschoben (Verschiebung Nr. 7)
2015 · 01 · 16 10:13

Martin Köchy

The International Geosphere Biosphere Program (IGBP) (Global Soil Data Task Group, 2000) produced a map of SOC stock on a 5' by 5' grid derived from the DSMW in conjunction with WISE data (v.1, 1125 profiles).

1.3.

Seite 13: [6] Gelöscht

Martin Köchy

2015 · 01 · 16 10:13

The World Reference Base Map of World Soil Resources (WRB, IUSS Working Group WRB, 2006), scale 1:25'000'000, is generalized from DSMW and includes updates from several databases not yet included in HWSD (v.1.2). The WRB contains 31 dominant soil type classes. Taxotransfer functions must yet be developed to derive organic C stocks from WRB.

The recently published Global Soil Dataset for Earth System Models (Shangguan et al., 2014) with a resolution of 0.5 arc minutes, uses information also used in the HWSD and additional information from the national databases of the USA, Canada, and Australia. Several harmonisation steps are applied to the data to derive amongst others soil carbon concentration, bulk soil density, and gravel content and depth for each soil mapping unit. The global organic carbon stocks after rasterizing the map to 0.5 arc minute pixels are reported as 1923, 1455 and 720 Pg for the upper 2.3, 1, and 0.3 m.

The accuracy of maps of soil C

1.4.

Seite 13: [7] Auf Seite 4 verschoben (Verschiebung Nr. 1)
2015-01-16 10:13

Martin Köchy

stocks depends on how well the soil mapping units are represented by soil profiles with complete characteristics. The latest ISRIC-WISE database (v.3.1) contains harmonized data of more than 10250 soil profiles (Batjes, 2009). The profiles, however, do not yet represent the terrestrial surface equally. Gaps include non-agricultural areas of North America, the Nordic countries, most parts of Asia (notably Iran, Kazakhstan, and Russia), Northern Africa, and Australia. To calculate SOC stocks one needs C_{org} , BD , soil depth, and volumetric gravel fraction.

1.5.

Seite 13: [8] Auf Seite 5 verschoben (Verschiebung Nr. 2)
2015-01-16 10:13

Martin Köchy

of the profiles (Batjes, 2009). BD and gravel fraction have low representation because they are seldom recorded during routine soil surveys. In numbers, 9970 profile descriptions include C_{org} in at least one layer, but of these only 3655 also include BD. Gravel fraction is explicitly indicated for 1100 of the 3655 profiles but earlier versions of the database could not distinguish between zero and absence of value. BD is included for 806 profiles where $C_{org} > 3\%$ and for 74 profiles where $C_{org} > 20\%$. The temporal origin of profile descriptions ranges from 1925 to 2005. The early data may no longer reflect current conditions where C input and decomposition rates may have changed. Efforts to expand the database of data-rich soil profiles and to use pedotransfer instead of taxotransfer functions has been going on since 1986 through the SOTER program (<http://www.isric.org/projects/soil-and-terrain-database-soter-programme>, accessed 2014-07-07, Nachtergaele, 1999).

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

| | | |
|-------------------------------------|--------------|----------------------|
| Seite 13: [9] Gelöscht organic C | Martin Köchy | 2015 · 01 · 16 10:13 |
|-------------------------------------|--------------|----------------------|

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 13: [9] Gelöscht Martin Köchy 2015 · 01 · 16 10:13
organic C

Seite 32: [10] Gelöscht Martin Köchy 2015 · 01 · 16 10:13

| Term | Definition |
|----------------------------|---|
| Concentration | organic carbon mass/soil dry mass, C_{org} |
| Content | organic carbon mass/soil volume = concentration \times bulk density |
| Areal density of fine soil | organic carbon mass/soil volume \times depth \times (1 – fractional volume of rocks, coarse roots, and ice) |
| Stock | areal density of fine soil integrated over all layers to a specified depth |
| Mass | stock integrated over a specified area |